

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) EP 0 739 886 A2

(12) EUROPEAN PATENT APPLICATION

(43) Date of publication:
30.10.1996 Bulletin 1996/44

(21) Application number: 96100433.0

(22) Date of filing: 12.01.1996

(51) Int. Cl.⁶: C07D 295/185, C07D 241/04,
C07C 257/18, C07C 257/22,
A61K 31/155, A61K 31/495

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE

(30) Priority: 28.04.1995 KR 9510383

(71) Applicant: LG Chemical Limited
Seoul (KR)


(72) Inventors:
• Oh, Yeong S.
Youseong-Ku, Daejeon (KR)
• Kim, Sang S.
Seo-ku, Daejeon (KR)
• Hwang, Sang Y.
Youseong-ku, Daejeon (KR)
• Yun, Mi K.
Seo-ku, Daejeon (KR)

- Hwang, Seong R.
Youseong-ku, Daejeon (KR)
- Hong, Seong W.
Daeduk-ku, Daejeon (KR)
- Lee, Yong H.
Youseong-ku, Daejeon (KR)
- Jeong, Yi N.
Youseong-ku, Daejeon (KR)
- Lee, Koo
Youseong-ku, Daejeon (KR)
- Shin, You S.
Youseong-ku, Daejeon (KR)

(74) Representative: KUHNEN, WACKER & PARTNER
Alois-Steinecker-Strasse 22
85354 Freising (DE)

(54) Selective thrombin inhibitors

(57) The present invention relates to a novel selective thrombin inhibitor having the following formula (I), which is also effective by oral administration:




in which

R¹ represents acetyl substituted with aryl or aryloxy, or represents sulfonyl substituted with substituted or unsubstituted aryl or N-containing heterocyclic group,

EP 0 739 886 A2

X represents a group of formula



- R² and R³ independently of one another represent hydrogen; cycloalkyl substituted or unsubstituted with carboxyl or alkoxycarbonyl; arylalkyloxy; hydroxy; or lower alkyl substituted or unsubstituted with carboxyl, alkoxycarbonyl or hydroxy, or
- R² and R³ together with nitrogen atom to which they are attached can form a piperidine group substituted with carboxyl or alkoxycarbonyl,
- R⁴ represents hydrogen, lower alkyl or lower alkoxy,
- R⁵ represents alkanesulfonyl; alkoxycarbonyl; alkylcarbonyl; formyl; lower alkyl; aryl substituted or unsubstituted with alkoxy or haloalkyl; or hydroxy-substituted lower alkyl, and
- R⁶ and R⁷ independently of one another represent hydrogen, lower alkyl or amino,


and to a process for preparation thereof and a pharmaceutical composition for thrombin inhibition which comprises the compound of formula (I) as an active ingredient.

Description

BACKGROUND OF THE INVENTION


1. Field of the Invention

The present invention relates to a novel selective thrombin inhibitor having the following formula (I):



in which

R^1 represents acetyl substituted with aryl or aryloxy, or represents sulfonyl substituted with substituted or unsubstituted aryl or N-containing heterocyclic group,
 X represents a group of formula



R^2 and R^3 independently of one another represent hydrogen; cycloalkyl substituted or unsubstituted with carboxyl or alkoxycarbonyl; arylalkyloxy; hydroxy; or lower alkyl substituted or unsubstituted with carboxyl, alkoxycarbonyl or hydroxy, or
 R^2 and R^3 together with nitrogen atom to which they are attached can form a piperidine group substituted with carboxyl or alkoxycarbonyl,
 R^4 represents hydrogen, lower alkyl or lower alkoxy,
 R^5 represents alkanesulfonyl; alkoxycarbonyl; alkylcarbonyl; formyl; lower alkyl; aryl substituted or unsubstituted with alkoxy or haloalkyl; or hydroxy-substituted lower alkyl, and
 R^6 and R^7 independently of one another represent hydrogen, lower alkyl or amino.

Some of the compounds of formula (I) can show an effective thrombin inhibitory activity even through oral administration and therefore, is very valuable.

The present invention also relates to a process for preparing the compound of formula (I) and to a pharmaceutical composition for thrombin inhibition which comprises the compound of formula (I) as an active ingredient.

2. Background Art

It has been generally known that the process for blood coagulation involves numerous complicated enzyme reactions of which the final step includes a reaction of converting prothrombin into thrombin. Thrombin produced from the final step of blood coagulation process activates platelets and converts fibrinogen into fibrin which is then converted into a higher molecular substance by polymerization and cross-linked by the action of activated blood factor XIII to form insoluble blood clotting. Accordingly, thrombin plays an important role in blood coagulation process. Thrombin also activates blood factors V and VIII which in turn accelerates the blood coagulation by a feed-back mechanism.

Thus, since thrombin inhibitors act as effective anti-coagulants and, at the same time, can inhibit the platelet activation and the production and stabilization of fibrin, for a long time, many attempts have been made to find out a method

10

25




35

4.



5

Further, Ro 46-6240 having the following formula (C) has been reported as the compound having an improved selectivity for thrombin over trypsin. This compound shows a possibility of development as an intravenous injectable formulation due to its short half-life in blood, but does not show any possibility for oral administration (see, J. Med. Chem. 1994, 37, 3889-3901).



(C)

In addition, a piperazine-based compound as recently developed has been reported as having somewhat possibility for oral administration in rat but having a low selectivity for thrombin (see WO 94/18185). Thus, such compounds did not come up to the expectation in this art.

Therefore, the present inventors have extensively studied to develop a certain compound which can be easily synthesized, shows an effective thrombin inhibitory activity with a high selectivity for thrombin over trypsin and can also be administered via oral route. As a result, we have identified that the thrombin inhibitor of formula (I) according to the present invention can achieve such purpose and thus completed the present invention.

Accordingly, it is an object of the present invention to provide a novel thrombin inhibitor of formula (I), as defined above, which can be administered via oral route and has a high selectivity for thrombin.


It is another object of the present invention to provide a process for preparing the thrombin inhibitor of formula (I).

Further, it is still another object of the present invention to provide a pharmaceutical composition for prevention of blood coagulation and treatment of various thrombosis, which comprises the thrombin inhibitor of formula (I) as an active ingredient.

The foregoing has outlined some of the more pertinent objects of the present invention. These objects should be construed to be merely illustrative of some of the more pertinent features and applications of the invention. Many other beneficial results can be obtained by applying the disclosed invention in a different manner or modifying the invention within the scope of the disclosure. Accordingly, other objects and a more thorough understanding of the invention may be had by referring to the disclosure of invention, in addition to the scope of the invention defined by the claims.

DISCLOSURE OF INVENTION

In one aspect, the present invention relates to a novel compound represented by the following formula (I):




(I)

its pharmaceutically acceptable salt, hydrate, solvate and isomer, in which

EP 0 739 886 A2

R^1 represents acetyl substituted with aryl or aryloxy, or represents sulfonyl substituted with substituted or unsubstituted aryl or N-containing heterocyclic group.
 X represents a group of formula



R^2 and R^3 independently of one another represent hydrogen; cycloalkyl substituted or unsubstituted with carboxyl or alkoxycarbonyl; arylalkyloxy; hydroxy; or lower alkyl substituted or unsubstituted with carboxyl, alkoxycarbonyl or hydroxy, or

R^2 and R^3 together with nitrogen atom to which they are attached can form a piperidine group substituted with carboxyl or alkoxycarbonyl,

R^4 represents hydrogen, lower alkyl or lower alkoxy,


R^5 represents alkanesulfonyl; alkoxycarbonyl; alkylcarbonyl; formyl; lower alkyl; aryl substituted or unsubstituted with alkoxy or haloalkyl; or hydroxy-substituted lower alkyl, and

R^6 and R^7 independently of one another represent hydrogen, lower alkyl or amino.

In the definition for each substituent of the compound of formula (I) according to the present invention, the term "lower alkyl" denotes a saturated, straight or branched hydrocarbon radical having 1 to 4 carbon atoms such as methyl, ethyl, isopropyl, isobutyl, t-butyl, etc.; the term "arylalkyloxy" denotes an alkoxy group substituted with aromatic ring such as benzyloxy, etc.; and the term "cycloalkyl" denotes a cyclic alkyl group having 3 to 8 carbon atoms such as cyclopentyl.

Among the compound of formula (I) above, the preferred one is the compounds in which

R^1 represents acetyl substituted with naphthyl or naphthyloxy, or represents sulfonyl substituted with naphthyl or phenyl which can be substituted or unsubstituted with one to four substituents selected from the group consisting of lower alkyl, lower alkoxy and dialkylamino,
 X represents a group of formula



R^2 and R^3 independently of one another represent C_{3-6} cycloalkyl substituted or unsubstituted with carboxyl or methoxycarbonyl; benzyloxy; lower alkyl substituted or unsubstituted with carboxyl, methoxycarbonyl or hydroxy; or hydroxy, or

R^2 and R^3 together with nitrogen atom to which they are attached can form a piperidine group substituted with carboxyl or methoxycarbonyl,

R^4 represents hydrogen,

R^5 represents methanesulfonyl, ethoxycarbonyl, formyl, ethyl, phenyl, methylcarbonyl, hydroxyethyl, or phenyl which can be substituted or unsubstituted with trifluoromethyl or ethoxy,

and

R^6 and R^7 independently of one another represent hydrogen, methyl or amino.

Typical example of the compound of formula (I) according to the present invention includes the following :

- (S)-N-cyclopentyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-butyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-propyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-(2-benzyloxyethyl)-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,

- (S)-N-cyclopentyl-N-butyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-ethyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-methyl-3-[4-(methylamidino)phenyl]-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-methyl-3-[4-(1,1-dimethylamidino)phenyl]-2-(2-naphthylsulfonylamino)propionamide,
- 5 • (S)-N-cyclopentyl-N-methyl-3-(4-amidrazonophenyl)-2-[(4-methoxy-2,3,6-trimethylbenzene)sulfonylamino]propionamide,
- (S)-N-cyclopentyl-N-hydroxy-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-(2-hydroxyethyl)-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-methyl-3-[4-(methylamidino)phenyl]-2-[(4-methoxy-2,3,6-trimethylbenzene)sulfonylamino]propionamide,
- 10 • (S)-N,N-dimethyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N,N-dimethyl-3-[4-(1-methylamidino)phenyl]-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclohexyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopropyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- 15 • (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(2-naphthalen-1-yl-acetyl-amino)-propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(5-dimethylamino-naphthalene-1-sulfonylamino)-propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(5-methoxy-naphthalene-1-sulfonylamino)-propionamide,
- (S)-2-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(6,7-dimethoxy-naphthalene-2-sulfonylamino)-propionamide,
- 20 • (S)-3-[4-(methylamidino)-phenyl]-N-cyclopentyl-N-methyl-2-(5-dimethylamino-naphthalene-1-sulfonylamino)-propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(naphthalene-1-sulfonylamino)-propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-[2-(naphthalene-1-yl-oxy)acetyl-amino]-propionamide,
- 25 • (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-[2-(naphthalene-2-yl-oxy)acetyl-amino]-propionamide,
- {[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino}-acetic acid methyl ester,
- {[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino}-acetic acid,
- (S)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid methyl ester,
- 30 • (S)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid,
- (R)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid methyl ester,
- (R)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid,
- (R)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-3-methyl-butyric acid methyl ester,
- 35 • (R)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-3-methyl butyric acid,
- 3-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid methyl ester,
- 40 • 3-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid,
- 4-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-butyric acid methyl ester,
- 4-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-butyric acid,
- [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopropylamino]-acetic acid methyl ester,
- 45 • [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopropylamino]-acetic acid,
- [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-butylamino]-acetic acid methyl ester,
- [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-butylamino]-acetic acid,
- [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopentylamino]-acetic acid methyl ester,
- 50 • [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopentylamino]-acetic acid,
- 1-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid methyl ester,
- 1-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid,
- 55 • 2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid ethyl ester,
- 2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid,

- (S)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-3-methyl-butiric acid methyl ester,
- 1-[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-piperidine-(R)-2-carboxylic acid methyl ester,
- 5 • 1-[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-piperidine-(R)-2-carboxylic acid,
- (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-(4-methylsulfonyl-piperazinyl)-2-oxoethyl]amide,
- (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-(4-ethoxycarbonyl-piperazinyl)-ethyl]amide,
- (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-(4-formyl-piperazinyl)-2-oxoethyl]amide,
- (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-(4-ethyl-piperazinyl)-2-oxoethyl]amide,
- 10 • (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-(4-phenyl-piperazinyl)-ethyl]amide,
- (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-[4-(3-trifluoromethylphenyl)-piperazinyl]-ethyl]amide,
- (S)-naphthalene-2-sulfonic acid [2-(4-acetyl-piperazinyl)-1-(4-amidrazono)benzyl-2-oxoethyl]amide,
- (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-[4-(2-hydroxyethyl)-piperazinyl]-ethyl]amide, and
- 15 • (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-[4-(2-ethoxyphenyl)-piperazinyl]-ethyl]amide.


The compound of formula (I) according to the present invention can also form a pharmaceutically acceptable salt. Suitable pharmaceutically acceptable salts for the compound (I) can include an acid addition salt formed with acids which can form a non-toxic acid addition salt containing pharmaceutically acceptable anion, for example, inorganic acids such as hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid, hydrobromic acid, hydroiodic acid, etc., organic carbonic acids such as tartaric acid, formic acid, citric acid, acetic acid, trichloroacetic acid, gluconic acid, benzoic acid, lactic acid, fumaric acid, maleic acid, etc., sulfonic acids such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, naphthalenesulfonic acid, etc., and the like acids.

In addition, since the compound of formula (I) according to the present invention can include an asymmetric carbon atom in its structure, it can be present in the form of racemate, diastereomeric mixture and individual diastereomer. All of such isomers are included within the scope of the present invention.

In another aspect, the present invention also relates to a process for preparing the compound of formula (I) as defined above.

According to the present invention, the compound of formula (I) can be prepared by reacting a compound of formula (II) with a compound of formula (III) as depicted in the following reaction scheme 1.

Reaction Scheme 1 :



In the above reaction scheme, X, R¹, R⁶ and R⁷ are defined as previously described.

As depicted in the above reaction scheme 1, the compound of formula (I) according to the present invention can be prepared by reacting the methylmercapto compound of formula (II) with the amine derivative of formula (III) as a nucleophilic substance. This reaction can be preferably carried out in the presence of a solvent. Although any organic solvent which does not adversely affect the reaction can be used in this reaction, in general, alcohol solvent such as methanol, ethanol, propanol, etc., is preferably used for this purpose.

In the above reaction, the reaction conditions including the amount of the reactants, reaction temperature, reaction time, etc. can be determined depending on the kind of particular reactant as used by a skilled person having an ordinary

EP 0 739 886 A2

knowledge in this art. Generally, although the reaction temperature can be varied within a substantial range, it is particularly preferable to carry out the reaction at 0°C to 50°C. In addition, the reaction generally takes 0.5 to 5 hours and can be preferably carried out for 1 to 2 hours.

After this reaction is completed, the reaction product can be separated and purified according to conventional working-up procedures, for example, chromatography, recrystallization, etc.

The methylmercapto compound of formula (II) used as the intermediate for preparing the compound of formula (I) in the reaction scheme 1 can be prepared according to the reaction scheme 2 or 3 as depicted below.

10

15

20

25

30


35

40


45

50


55

Reaction Scheme 2 :


[1]




[2]




[3]




[4]



[5]



(II)

Reaction Scheme 3:

In the above reaction schemes,

55 X, R¹, R⁶ and R⁷ are defined as previously described, and
 P represents an amino-protecting group.

Hereinafter, the reaction schemes 2 and 3 are specifically explained.

In the reaction scheme 2, first the C-terminal of the compound [1] is coupled with the amine group X to prepare the compound [2] from which the amino-protecting group at N-terminal is removed to prepare the compound [3]. Then, the group R¹ is introduced into the deprotected N-terminal of the compound [3] to prepare the compound [4]. Alternatively, according to the reaction scheme 3, first the group R¹ is introduced into the N-terminal of the compound [7] and then the C-terminal is coupled with the amine group X to prepare the compound [4].

Specifically, according to the reaction scheme 2, the compound of formula [1] is coupled with the amine compound corresponding to the substituent X to obtain the compound of formula [2] from which the amino-protecting group at the N-terminal is removed to obtain the compound of formula [3]. Then, the group R¹ is introduced into the N-terminal of the compound [3] to produce the nitrile compound of formula [4] which is then saturated with hydrogen sulfide in the presence of pyridine and triethylamine to produce the thioamide compound of formula [5]. This thioamide compound is then methylated with a methylating agent such as iodomethane, dimethylsulfate, methyltriflate, etc., to obtain the desired methylmercapto compound of formula (II).

According to the reaction scheme 3, first the group R¹ is introduced into the N-terminal of the compound of formula [7] by reacting the compound [7] with the compound of formula [6] and then the amine group is introduced into the C-terminal of the resulting compound of formula [8] by coupling the compound [8] with the amine compound corresponding to the substituent X to obtain the compound of formula [4] which is then subjected to the same procedure as in the reaction scheme 2 to produce the desired methylmercapto compound of formula (II).

The coupling agent which can be used for the coupling procedure in the reaction schemes 2 and 3 includes one or more substances selected from the group consisting of dicyclohexylcarbodiimide(DCC), 3-ethyl-3'-(dimethylamino)propylcarbodiimide(EDC), bis-(2-oxo-3-oxazolidinyl)phosphinic acid chloride(BOP-Cl) and diphenylphosphorylazide(DPPA) but is not limited thereto.

Although the carboxylic acid compounds [1] and [7] used in the reaction schemes 2 and 3 can be used in their free acid forms, they can be preferably used in the form of their reactive derivatives, for example, an acid halide derivative or other activated ester derivative to facilitate the reaction. Particularly, the activated ester derivative of carboxylic acid is necessary for a coupling reaction with an amine compound to form an amide linkage or a coupling reaction with an alcohol to form an ester linkage. Such reactive derivatives include conventional derivatives which can be prepared according to the method conventionally used in this technical field. For example, as the acid halide derivative an acid chloride is included; and the activated ester derivative includes a carboxylic acid anhydride derived from alkoxycarbonyl halide such as methoxycarbonylchloride, isobutyloxycarbonylchloride, etc., and a coupling agent, N-hydroxyphthalimide-derived ester, N-hydroxysuccinimide-derived ester, N-hydroxy-5-norbornene-2',3'-dicarboxyimide-derived ester, 2,4,5-trichlorophenol-derived ester, etc., but is not limited thereto.

The thrombin inhibitory effect of the compound of formula (I) according to the present invention can be identified by determination of a dissociation constant K_i represented by the following equation according to the known method described in literature [see, Methods in enzymology V. 80, p341-361; Biochemistry 27, p2144-2151 (1988)].

$$K_i = \frac{[E] \cdot [I]**}{[EI]**}$$

* [E] : concentration of the free enzyme
 ** [I] : concentration of the unbound inhibitor
 *** [EI] : concentration of the enzyme-inhibitor complex

The dissociation constant K_i denotes the degree of dissociation of the enzyme-thrombin inhibitor complex. Accordingly, the low dissociation constant means the high binding property of thrombin inhibitor to enzyme and therefore, is estimated that the thrombin inhibitor has a high inhibitory activity for thrombin. Such dissociation constant can be determined by reacting thrombin with a certain substrate which develops a color when it is hydrolyzed by the action of thrombin, and then measuring the degree of color development as a function of time by means of spectrophotometry.

In the present invention, Chromozym TH (Gly-Pro-Arg-4-nitroanilide acetate) is used as the substrate substance for thrombin, which develops the color by the action of thrombin. Chromozym TH is hydrolyzed by thrombin to produce yellow para-nitroaniline. Accordingly, the amount of yellow para-nitroaniline thus produced can be measured as the change of absorbance in the course of time to determine the thrombin inhibitory activity of the compound according to the present invention. That is, the enzyme activity can be determined from the rate of change in absorbance and then can be directly connected with the ability of the thrombin inhibitor to inhibit the enzyme activity (see, Methods in Enzymology, V. 80, p341-361, Biochemistry 27, p2144-2151, 1988).

To identify the selectivity for thrombin over trypsin of the compound of the present invention, the inhibitory activity for trypsin of the compound of formula (I) is measured as K_i value according to the same method as the above method for determination of thrombin inhibitory activity, and then the ratio of the activity for trypsin to the activity for thrombin is calculated. In this case, the procedure for determination of inhibitory activity for trypsin is substantially identical to that for thrombin, except that N-benzoyl-Val-Gly-Arg para-nitroanilide hydrochloride is used as the substrate.

As the result of determination of inhibitory activity for thrombin and trypsin of the compound of formula (I) according to the present invention, it can be identified that the compound of the present invention shows an excellent thrombin inhibitory activity and further has a high selectivity for thrombin against trypsin. Particularly, the selectivities of compounds of Examples 1 and 7 for thrombin against trypsin are approximately 2900 times and 26304 times, respectively, whereas the selectivities of the known thrombin inhibitors, argatroban (A) and NAPAP (B), are merely 250 times and 50 times, respectively. Accordingly, it can be seen that the compound (I) of the present invention achieves a significant improvement in the selectivity for thrombin against trypsin.

As mentioned above, since the novel compound of formula (I) according to the present invention is a thrombin inhibitor which shows a potent thrombin inhibitory activity even through oral administration and also has a high selectivity for thrombin over trypsin, it is useful for prophylaxis of blood coagulation and treatment of various thrombosis.

Accordingly, it is the third object of the present invention to provide a pharmaceutical composition for prophylaxis of blood coagulation and treatment of thrombosis which comprises the compound of formula (I) or its pharmaceutically acceptable salt as an active ingredient.

When the compound of formula (I) according to the present invention is administered to the subject host for clinical purpose, a daily dosage of the compound (I) can be varied preferably in the range of 0.001mg to 10mg per kg of body weight. However, if required for a certain patient, a specific dosage beyond the above mentioned dosage range can be determined depending on the a specific compound to be used, weight, sex and healthy condition of the respective patient, diet, administration time and method, secretion rate of the compound, other active component to be combined and severity of the diseases to be treated.

The compound of the present invention can be administered in the form of an injectable preparation or an oral preparation according to the desired purpose.

The injectable preparation, for example, sterilized, injectable aqueous or oily suspensions can be formulated using a suitable dispersing agent, wetting agent or suspending agent according to the known method conventionally used in the field of preparation of injections. As the aqueous solvent suitable for this purpose, water, Ringer's solution or isotonic NaCl solution can be used. In addition, although a sterilized fixing oil can be used as the solvent or suspending medium, any non-irritable fixing oil including mono- or di-glycerides can be used for the same purpose. Furthermore, a fatty acid such as oleic acid may be added to the injectable preparation.

The solid preparation for oral administration can be in the form of capsules, tablets, pills, powders and granules, with the capsule or tablet formulation being particularly useful. The tablets and pills can be prepared preferably in the enteric-coated formulation. In formulating the solid preparation, the active compound of formula (I) according to the present invention can be combined with a pharmaceutically acceptable carrier, for example, one or more inert diluent(s) such as sucrose, lactose, starch, etc., lubricants such as magnesium silicate, disintegrants, binders and the like.

One of the major characteristic features of the compound of formula (I) according to the present invention is that the compound of formula (I) exhibits a good pharmacological effect even when it is formulated in the form of an oral preparation and then administered via oral route. This can be demonstrated from the result of pharmacokinetic experiments in rat and dog as the experimental animal. That is, in such experiments it could be identified that the active compound of the present invention is retained in blood for a prolonged period when it is administered via oral route. Accordingly, the compound of formula (I) according to the present invention is more useful in view of the fact that it can be effectively used in the form of oral preparation.

Further, from the pharmacokinetic experiments it could also be identified that the active compound of the present invention can achieve the desired purpose without acute toxicity in mammals including rat and dog.

When the thrombin inhibitor according to the present invention is administered to obtain anti-coagulant and thrombolytic effects, it can be administered in combination with one or more substance(s) selected from the group consisting of thrombolytic agents and agents for inhibition of thrombocyte activity. As the thrombolytic agent which can be used for this purpose, t-PA, urokinase, streptokinase, etc. can be mentioned; and aspirin, ticlopidin, clopidogrel, 7E3 monoclonal antibody, etc. can be used as the agent for inhibition of thrombocyte activity.

However, it should be understood that the preparation containing the active compound of the present invention for treatment and prophylaxis of thrombosis is not limited to those described above and can include any preparation useful for the same purpose.

Hereinafter, the present invention will be more specifically explained by the working examples. However, it should be understood that the examples are given only for the explanation of the present invention and not intended to limit the present invention in any manner.

Preparation 1

Synthesis of cyclopentyl-methylamine

To a solution of cyclopentanone(10ml, 113 mmol) in methanol (50ml) and water(50ml) were added methylamine hydrochloride(7.6g, 113 mmol) and sodium cyanoborohydride (NaBH_3CN) (7.1g, 113 mmol). The mixture was heated

EP 0 739 886 A2

to reflux for 12hrs at pH 6. Methanol was evaporated under reduced pressure and the residue was cooled to 0°C, adjusted to pH 2 using 3N hydrochloric acid and then washed three times with diethyl ether. The aqueous layer was cooled again to 0°C and then adjusted to pH 11 using 6N sodium hydroxide solution. To this mixture was added t-butyloxycarbonyl anhydride (24.5g, 113 mmol) in dioxane(50 ml). The solution was stirred for 3 hours at room temperature and concentrated under reduced pressure to about 30ml. The residue was extracted with ethyl acetate and washed with aqueous 0.5N hydrochloric acid and saturated sodium bicarbonate solutions. The organic layer was dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure to obtain the white solid which was then purified by column chromatography(eluent = ethyl acetate:hexane = 7:3 (v/v)). Upon purification, the obtained solid product was dissolved in 4N HCl-dioxane solution(60ml) and the resulting solution was stirred for 30 minutes at room temperature. The solvent was evaporated in vacuo to obtain the title compound (13.7g Yield :90.5%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 3.50(m, 1H), 2.68(s, 3H), 2.10(m, 2H), 1.86-1.50(m, 6H)

Preparation 2

Synthesis of (S)-N-cyclopentyl-N-methyl-3-(4-cyanophenyl)-2-(butyloxycarbonylamino)propionamide

To a solution of (S)-3-(4-cyanophenyl)-2-(butyloxycarbonylamino)propionic acid (0.7g, 2.41 mmole) in dimethylformamide (DMF, 6ml) were added, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride(EDC, 0.7g) and 1-hydroxybenzotriazole hydrate(HOBT, 0.4g) at 0°C. The mixture was stirred until they are completely dissolved therein. To this reaction mixture were added the compound (0.4g, 2.96 mmole) prepared in Preparation 1 and N-methylmorpholine(1.0 ml) and then the reaction temperature was slowly elevated to room temperature. The reaction solution was stirred for 3.5 hours. After the reaction is completed, the reaction solution was concentrated under reduced pressure to remove the volatile substances and the remaining residue was diluted with ethyl acetate, washed successively with aqueous saturated sodium hydrogen carbonate solution, dilute hydrochloric acid and brine, dried over anhydrous sodium sulfate, filtered and then concentrated. The residue was purified by column chromatography (ethyl acetate : hexane = 7:3 (v/v)) to obtain the purified title compound (0.65g, Yield : 73.0%).

$^1\text{H NMR}(\text{CDCl}_3, \text{ppm}) \delta$: 7.61(m, 2H), 7.32(m, 2H), 5.48, 5.01-4.86, 4.12(3m, 3H), 2.75, 2.62(2s, 3H), 2.90-1.20(m, 17H)

Mass(FAB, m/e) : 372($\text{M}^+ + 1$)

Preparation 3

Synthesis of (S)-N-cyclopentyl-N-methyl-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino)propionamide

The compound (0.65g, 1.75 mmole) prepared in Preparation 2 was dissolved in dichloromethane(3ml) and then cooled to -10°C, and trifluoroacetic acid(TFA, 1ml) was added thereto. The reaction mixture was stirred for 5 minutes, slowly warmed to room temperature, stirred for 30 minutes and then concentrated under reduced pressure to remove the volatile substances. The residue was dried by means of a vacuum pump and then 6ml of DMF was added thereto. The mixture was cooled to 0°C and of N,N-diisopropylethylamine(1ml) was added thereto. The reaction mixture was warmed to room temperature and stirred for about 5 minutes. After addition of 2-naphthalenesulfonyl chloride (0.47g, 2.07 mmole), the reaction mixture was stirred for one hour to complete the reaction and then concentrated under reduced pressure to remove the volatile substances. The residue was diluted with ethyl acetate, washed with saturated sodium hydrogen carbonate solution and brine, dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by column chromatography (ethyl acetate : hexane = 1:1 (v/v)) to obtain the title compound (0.65g, Yield : 80.2%).

$^1\text{H NMR}(\text{CDCl}_3, \text{ppm}) \delta$: 8.28(m, 1H), 7.87(m, 3H), 7.73(m, 3H), 7.49(m, 2H), 5.92(m, 1H), 4.50, 4.32, 3.76(m, m, 2H), 2.95(m, 2H), 2.36, 2.22(s, s, 3H), 1.60-1.20(m, 6H), 0.98, 0.80, 0.47(m, m, 2H)

Mass(FAB, m/e) : 462($\text{M}^+ + 1$)

Example 1

Synthesis of (S)-N-cyclopentyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The compound (0.65g, 1.41 mmole) prepared in Preparation 3 was dissolved in of pyridine(10ml) and the resulting solution was introduced into a branched flask, to which triethylamine(0.45ml) was added. The reaction flask was

equipped so that hydrogen sulfide(H_2S) gas can be slowly introduced through one branch of the flask and discharged through another branch. The reaction solution was saturated with hydrogen sulfide gas, while stirring for about 10 minutes, during which the colorless solution was changed into green color and then gradually into dark brown. The flask was closed with rubber stopper and allowed to stand for 3 days at room temperature to complete the reaction. Then, the reaction solution was distilled under reduced pressure to remove the volatile substances and dried by means of a vacuum pump. To the obtained yellow solid were added acetone(15ml) and iodomethane(CH_3I , 0.65ml) together and this mixture was heated to reflux for 30 minutes. This reaction mixture was distilled again under reduced pressure to remove the volatile substances and dried by means of a vacuum pump. The residue was dissolved in absolute methanol(8ml) and then stirred. To this mixture was added portionwise 80% hydrazine hydrate ($H_2NNH_2 \cdot H_2O$, 0.12ml, 1.98 mmole) over three times at an interval of 10 minutes. After the reaction is completed, the reaction solution was concentrated and then purified by HPLC to obtain the title compound (0.63g, Yield : 73.0%).

HPLC condition :

○ eluent = methanol : water (75:25 v/v), each of which contains CF_3COOH in the concentration of 0.1%

○ : wavelength = 215nm

○ : elution rate = 20ml/min.

○ column = Delta PAK C_{18} 100Å (30x300mm)

1H NMR(CD_3OD , ppm) δ : 8.28(d, 1H), 7.92(m, 3H), 7.70-7.50(m, 5H), 7.35(dd, 2H), 4.60, 4.45(t, t, 1H), 4.12, 3.99(m, m, 1H), 3.00, 2.87(m, m, 2H), 2.49, 2.26(s, s, 3H), 1.60-1.00, 0.73-0.52(m, m, 8H)

Mass(FAB, m/e) : 494($M^+ + 1$)

Preparation 4

Synthesis of butyl-methylamine trifluoroacetic acid salt

N-butyloxycarbonylbutylamine(140mg, 0.80 mmole) was dissolved in DMF (8ml), and sodium hydride(NaH , 20mg, 1 eq. wt.) and iodomethane(0.10ml, 2 eq. wt.) were added thereto. The reaction mixture was stirred for 30 minutes at room temperature and filtered through celite bed and then concentrated under reduced pressure to remove the solvent. The residue was diluted with ethyl acetate, washed with 0.5N aqueous hydrochloric acid solution, dried over anhydrous magnesium sulfate and filtered. The organic layer was concentrated under reduced pressure and dried by means of a vacuum pump to obtain a white solid, which was then dissolved in dichloromethane and cooled down to 0°C. To this mixture was added 1ml of trifluoroacetic acid(TFA). The reaction solution was stirred for 30 minutes at room temperature, concentrated under reduced pressure and then dried by means of a vacuum pump to obtain the title compound(0.16g) in a quantitative yield.

1H NMR($CDCl_3$, ppm) δ : 1.02(t, 3H), 1.30-1.80(m, 4H), 3.04(s, 3H), 3.52(t, 2H), 8.20(s, 2H)

Example 2

Synthesis of (S)-N-butyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The compound prepared in Preparation 4 was reacted according to the same procedure as Preparations 2 and 3 to obtain the intermediate (S)-N-butyl-N-methyl-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino) propionamide(0.23g) which was used as the starting material. This starting material was treated according to the same procedure as Example 1 to obtain the purified title compound(0.1g, Yield : 40.0%).

1H NMR(CD_3OD , ppm) δ : 8.30(d, 1H), 7.98(m, 3H), 7.81-7.30(m, 7H), 4.50(m, 1H), 3.25-2.55(m, 4H), 2.78, 2.45(2s, 3H), 1.40-0.50(m, 7H)

Mass(FAB, m/e) : 482($M^+ + 1$)

Example 3

Synthesis of (S)-N-cyclopentyl-N-propyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The same procedure as Preparation 1, except that propylamine was used instead of methylamine, was carried out to obtain cyclopentyl-propylamine hydrochloride which was then treated according to the same procedure as Preparations 2 and 3 to obtain the intermediate (S)-N-cyclopentyl-N-propyl-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino)pro-

EP 0 739 886 A2

pionamide(0.15g). This intermediate compound was used as the starting material and treated according to the same procedure as Example 1 to obtain the purified title compound(0.089g, Yield : 55.1%).

¹H NMR(CD₃OD, ppm) δ : 8.35-7.35(m, 11H), 4.62-4.32(m, m, 1H), 3.90, 3.70(m, m, 1H), 3.10-2.50(m, 4H),
1.70-0.50(m, 13H)
Mass(FAB, m/e) : 522(M⁺+1)

Example 4

Synthesis of (S)-N-cyclopentyl-N-(2-benzyloxyethyl)-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The same procedure as Preparation 1, except that 2-benzyloxyethylamine was used instead of methylamine, was carried out to obtain cyclopentyl-(2-benzyloxyethyl)amine hydrochloride which was then treated according to the same procedure as Preparations 2 and 3 to obtain the intermediate (S)-N-cyclopentyl-N-(2-benzyloxyethyl)-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino) propionamide(0.15g). This intermediate compound was used as the starting material and treated according to the same procedure as Example 1 to obtain the purified title compound(0.15g, Yield : 93.8%).

¹H NMR(CD₃OD, ppm) δ : 8.30-7.15(m, 16H), 4.64(m, 1H), 4.48, 4.39(s, s, 1H), 4.14, 4.00, 3.75, 3.45 (m, m, m, 3H), 3.10-2.70(m, 5H), 1.62-1.00(m, 8H)
Mass(FAB, m/e) : 614(M⁺+1)

Example 5

Synthesis of (S)-N-cyclopentyl-N-butyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The same procedure as Preparation 1, except that butylamine was used instead of methylamine, was carried out to obtain butylcyclopentylamine hydrochloride which was then treated according to the same procedure as Preparations 2 and 3 to obtain the intermediate (S)-N-cyclopentyl-N-butyl-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino)propionamide(0.28g). This intermediate compound was used as the starting material and treated according to the same procedure as Example 1 to obtain the purified title compound(0.18g, Yield : 60%).

¹H NMR(CD₃OD, ppm) δ : 8.32(d, 1H), 7.96(m, 3H), 7.78-7.55(m, 5H), 7.42(dd, 2H), 4.62, 4.30(m, m, 1H), 4.02, 3.90(m, m, 1H), 3.10-2.75, 2.55(m, m, 4H), 1.65-0.80(m, 12H), 0.65 (t, 3H)
Mass(FAB, m/e) : 536(M⁺+1)

Example 6

Synthesis of (S)-N-cyclopentyl-N-ethyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The same procedure as Preparation 1, except that ethylamine was used instead of methylamine, was carried out to obtain cyclopentyl-ethylamine hydrochloride which was then treated according to the same procedure as Preparations 2 and 3 to obtain the intermediate (S)-N-cyclopentyl-N-ethyl-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino)propionamide(0.21g). This intermediate compound was used as the starting material and treated according to the same procedure as Example 1 to obtain the purified title compound(0.11g, Yield : 50.0%).

¹H NMR(CD₃OD, ppm) δ : 8.32(d, 1H), 7.97(m, 3H), 7.75-7.55(m, 5H), 7.42(m, 2H), 4.60, 4.38(m, m, 1H), 3.98, 3.87(m, m, 1H), 3.20-2.70(m, 4H), 1.65-1.00(m, 8H), 0.95, 0.58(t, t, 3H)
Mass(FAB, m/e) : 508(M⁺+1)

Example 7

Synthesis of (S)-N-cyclopentyl-N-methyl-3-[4-(methanimidino)-phenyl]-2-(2-naphthylsulfonylamino)propionamide

The intermediate compound prepared in Preparation 3 was used as the starting material and treated according to the same procedure as Example 1, except that methylamine, instead of hydrazine, was added over 3 times at an interval of one hour, to obtain the purified title compound(0.064g, Yield : 8%).

EP 0 739 886 A2

^1H NMR(CD_3OD , ppm) δ : 0.50-1.60(m, 8H), 2.00-2.49(2s, 3H), 2.84(m, 1H), 2.96(m, 1H), 3.00(s, 3H), 4.05(m, 1H), 4.50(m, 1H), 7.20-8.30(m, 11H)

Mass(FAB, m/e) : 493($\text{M}^+ + 1$)

5 Example 8

Synthesis of (S)-N-cyclopentyl-N-methyl-3-[4-(1,1-dimethylamidino)phenyl]-2-(2-naphthylsulfonylamino)propionamide

The intermediate compound prepared in Preparation 3 was used as the starting material and treated according to the same procedure as Example 1, except that dimethylamine, instead of hydrazine, was added over three times at an interval of one hour, to obtain the purified title compound(0.18g, Yield : 22%).

^1H NMR(CD_3OD , ppm) δ : 0.60-1.60(m, 8H), 2.29, 2.54(2s, 3H), 2.95(m, 1H), 3.06(m, 1H), 3.09(s, 3H), 3.31(s, 3H), 4.16(m, 1H), 4.60(m, 1H), 7.20-8.30(m, 11H)

15 Mass(FAB, m/e) : 507($\text{M}^+ + 1$)

Example 9

Synthesis of (S)-N-cyclopentyl-N-methyl-3-(4-amidrazonophenyl)-2-(4-methoxy-2,3,6-trimethylbenzenesulfonylamino)propionamide

The compound prepared in Preparation 2 was treated according to the same procedure as Preparation 3, except that 4-methoxy-2,3,6-trimethylbenzenesulfonyl chloride was used instead of 2-naphthalenesulfonyl chloride, to obtain the intermediate (S)-N-cyclopentyl-N-methyl-3-(4-cyanophenyl)-2-(4-methoxy-2,3,6-trimethylbenzenesulfonylamino)propionamide(0.27g). This intermediate compound was used as the starting material and treated according to the same procedure as Example 1 to obtain the purified title compound(0.16g, Yield : 57.1%).

^1H NMR(CD_3OD , ppm) δ : 7.62(d, 2H), 7.40(dd, 2H), 6.70(d, 1H), 4.35, 3.90(m, m, 2H), 3.83(d, 3H), 3.00 (m, 2H), 2.50(m, 9H), 2.1(s, 3H), 1.75-0.90(m, 8H)

30 Mass(FAB, m/e) : 516($\text{M}^+ + 1$)

Preparation 5

Synthesis of cyclopentyl-hydroxylamine

Hydroxylamine hydrochloride($\text{H}_2\text{NOH} \cdot \text{HCl}$, 5.0g, 71.95 mmole) was dissolved in water(14ml), and methanol(30ml) and cyclopentanone (5.1ml, 57.66 mmole) were added thereto. The mixture was stirred, cooled to 0°C and then adjusted to pH 8 by adding 6N aqueous sodium hydroxide solution. Sodium cyanoborohydride (NaBH_3CN , 1.9g, 30.24 mmole) was added thereto. The mixture was warmed to room temperature and then stirred. The reaction mixture was retained at pH 4 by adding portionwise the mixed solution of 6N HCl(20ml) and methanol(30ml) in the course of the reaction. After 5 hours, the reaction mixture was adjusted to pH 7 and distilled under reduced pressure to remove methanol. The remaining reaction solution was cooled down to 0°C , adjusted again to pH 11, saturated with sodium chloride and then extracted four times with chloroform. The extract was dried over anhydrous magnesium sulfate, filtered and concentrated to obtain the title compound (3.4g, Yield : 58.3%).

45 ^1H NMR(CDCl_3 , ppm) δ : 7.20-5.00(bs, 1H), 3.56(m, 1H), 1.90-1.45(m, 9H)

Example 10

Synthesis of (S)-N-cyclopentyl-N-hydroxy-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The compound prepared in Preparation 5 was treated according to the same procedure as Preparations 2 and 3 to obtain the intermediate (S)-N-cyclopentyl-N-hydroxy-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino)propionamide(0.1g). This intermediate compound was used as the starting material and treated according to the same procedure as Example 1 to obtain the purified title compound (0.07g, Yield : 63.6%).

^1H NMR(CD_3OD , ppm) δ : 8.25-7.30(m, 11H), 4.75(m, 1H), 4.30(m, 1H), 3.10, 2.75(m, m, 2H), 1.70-1.00(m, 8H)

Mass(FAB, m/e) : 496($\text{M}^+ + 1$)

EP 0 739 886 A2

Example 11

Synthesis of (S)-N-cyclopentyl-N-(2-hydroxyethyl)-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The compound (0.15g, 0.24 mmole) prepared in Example 4 was dissolved in methanol (10ml) and then palladium hydroxide (0.02g) was added thereto. Hydrogen balloon was attached to the reaction vessel. After stirring for 2 days, the reaction mixture was filtered through a celite bed and concentrated. The residue was purified by HPLC to obtain the title compound (0.08g, Yield : 63.7%). HPLC condition was identical to that used in Example 1.

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 7.80-7.00(m, 11H), 4.47(m, 1H), 4.00(m, 1H), 3.60(m, 2H), 3.10-2.70(m, 4H), 1.70-1.10(m, 8H)
Mass(FAB, m/e) : 524($\text{M}^+ + 1$)

Example 12

Synthesis of (S)-N-cyclopentyl-N-methyl-3-[4-(methylamidino)-phenyl]-2-(4-methoxy-2,3,6-trimethylbenzenesulfonylamino)propionamide

The same procedure as Example 9, except that methylamine, instead of hydrazine, was added over three times at an interval of one hour, was carried out to obtain the purified title compound (0.08g, Yield : 28.3%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 7.65(d, 2H), 7.40(dd, 2H), 6.70(d, 1H), 4.40, 3.90(m, m, 2H), 3.85(d, 3H), 3.20 - 2.90(m, 5H), 2.55(m, 9H), 2.10(s, 3H), 1.75-0.90(m, 8H)
Mass(FAB, m/e) : 515($\text{M}^+ + 1$)

Example 13

Synthesis of (S)-N,N-dimethyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The same procedure as Preparations 2 and 3, except that dimethylamine was used, was carried out to obtain the intermediate (S)-N,N-dimethyl-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino)-propionamide (0.11g). This intermediate compound was used as the starting material and treated according to the same procedure as Example 1 to obtain the purified title compound (0.080g, Yield : 53%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 2.19(s, 3H), 2.62(s, 3H), 2.70-2.95(m, 2H), 4.41(m, 1H), 7.20-8.30(m, 11H)
Mass(FAB, m/e) : 440($\text{M}^+ + 1$)

Example 14

Synthesis of (S)-N,N-dimethyl-3-[4-(methylamidino)phenyl]-2-(2-naphthylsulfonylamino)propionamide

The same procedure as Example 13, except that instead of hydrazine in the procedure of Example 1, methylamine is added over three times at an interval of one hour, was carried out to obtain the purified title compound (0.10g, Yield : 48%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 2.31(s, 3H), 2.75(s, 3H), 2.80-3.05(m, 2H), 3.07(s, 3H), 4.54(m, 1H), 7.30-8.40(m, 11H)
Mass(FAB, m/e) : 439($\text{M}^+ + 1$)

Example 15

Synthesis of (S)-N-cyclohexyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The same procedure as Preparation 1, except that cyclohexanone was used instead of cyclopentanone, was carried out to obtain cyclohexyl-methylamine hydrochloride, which was then treated according to the same procedure as Preparations 2 and 3 to obtain the intermediate (S)-N-cyclohexyl-N-methyl-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino)propionamide (0.21g). This intermediate compound was used as the starting material and treated according to the same procedure as Example 1 to obtain the purified title compound (0.17g, Yield : 73.9%).

EP 0 739 886 A2

¹H NMR(CD₃OD, ppm) δ : 8.32(d, 1H), 7.95(m, 3H), 7.75-7.53(m, 5H), 7.40(dd, 2H), 4.50, 4.21, 3.62(m, m, m, 2H), 3.05(m, 1H), 2.90(m, 1H), 2.55, 2.40(s, s, 3H), 1.80-0.62(m, 10H)

Mass(FAB, m/e) : 508(M⁺+1)

5 Example 16

Synthesis of (S)-N-cyclopropyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide

The same procedure as Preparation 1, except that cyclopropanone was used instead of cyclopentanone, was carried out to obtain cyclopropyl-methylamine hydrochloride, which was then treated according to the same procedure as Preparations 2 and 3 to obtain the intermediate (S)-N-cyclopropyl-N-methyl-3-(4-cyanophenyl)-2-(2-naphthylsulfonylamino)propionamide(0.21g). This intermediate compound was used as the starting material and treated according to the same procedure as Example 1 to obtain the purified title compound(0.06g, Yield : 12%).

¹H NMR(CD₃OD, ppm) δ : 0.40-0.90(m, 4H), 2.40(s, 3H), 2.75(m, 1H), 2.95(m, 1H), 7.20-8.30(m, 11H)
Mass(FAB, m/e) : 466(M⁺+1)

Preparation 6

20 Synthesis of (S)-3-(4-cyanophenyl)-N-cyclopentyl-N-methyl-2-(2-naphthalen-1-yl-acetyl-amino)-propionamide

The compound(0.51g, 1.34 mmole) prepared in Preparation 2 was dissolved in 3ml of dichloromethane and cooled down to -10°C, and then trifluoroacetic acid(TFA, 3ml) was added thereto. The reaction mixture was stirred for 5 minutes, slowly warmed to room temperature, stirred again for 30 minutes and then distilled under reduced pressure to remove the volatile substance. The residue was dried using a vacuum pump and then DMF(10ml) was added thereto. This solution was cooled to -10°C, and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride(EDC, 0.4g) and 1-hydroxybenzotriazole(HOBT, 0.2g) were added thereto and then stirred until they are completely dissolved. To the resulting solution were added 1-naphthalene acetic acid(0.26g, 1.4 mmole) and N,N-diisopropylethylamine(1.2ml). The reaction mixture was slowly warmed to room temperature and stirred for 3 hours. Upon completion of the reaction, the reaction solution was distilled under reduced pressure to remove the volatile substance. The residue was diluted with ethyl acetate, washed successively with aqueous saturated sodium hydrogen carbonate solution, dilute hydrochloric acid and brine, dried over anhydrous magnesium sulfate and then filtered. The filtrate was concentrated and the residue was purified by column chromatography(methanol:chloroform=1:99 (v/v)) to obtain the title compound (0.42g, Yield : 68%).

¹H NMR(CDCl₃, ppm) δ : 8.0-6.8(m, 11H), 6.5(m, 1H), 5.1-5.3(m, 1H), 4.8-4.2(m, 1H), 4.1-3.8(m, 2H), 3.0-2.6(m, 5H), 1.8-1.3(m, 8H)

Example 17

40 Synthesis of (S)-3-[4-(amidrazono)phenyl]-N-cyclopentyl-N-methyl-2-(2-naphthalen-1-yl-acetyl-amino)propionamide

The compound(0.21g, 0.48 mmole) prepared in Preparation 6 was dissolved in pyridine(3ml) and the resulting solution was introduced into a branched flask, to which triethylamine(0.2ml) was added. The reaction flask was equipped so that hydrogen sulfide(H₂S) gas can be slowly introduced through one branch of the flask and discharged through another branch. The reaction solution was saturated with hydrogen sulfide gas while stirring for about 10 minutes, during which the colorless solution was changed into green color and then gradually into dark brown. The flask was closed with rubber stopper and allowed to stand for 3 days at room temperature to complete the reaction. Then, the reaction solution was distilled under reduced pressure to remove the volatile substance and dried by means of a vacuum pump. To the obtained yellow solid were added acetone(10ml) and iodomethane(CH₃I, 0.3ml) together and the mixture was heated under reflux for 30 minutes. This reaction mixture was distilled again under reduced pressure to remove the volatile substance and dried by means of a vacuum pump. The residue was dissolved in absolute methanol(5ml) and then stirred. To this mixture was added portionwise 80% hydrazine hydrate (H₂NNH₂ · H₂O, 0.04ml, 0.72 mmole) over three times at an interval of 10 minutes. After the reaction is completed, the reaction solution was concentrated and then purified by HPLC to obtain the title compound(0.16g, Yield : 70%).

¹H NMR(CD₃OD, ppm) δ : 8.0-7.3(m, 11H), 5.3-5.1(m, 1H), 4.8-4.2(m, 1H), 3.9(s, 2H), 3.2-2.9(m, 2H), 2.78, 2.72(2s, 3H), 1.8-1.3(m, 8H)
Mass(FAB, m/e) : 472(M⁺+1)

Example 18Synthesis of (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(5-dimethylamino-naphthalene-1-sulfonylamino)-propionamide

The same procedure as Preparation 3, except that 5-(N,N-dimethylamino)-1-naphthalenesulfonyl chloride was used instead of 2-naphthalenesulfonyl chloride, was carried out to obtain the intermediate (S)-3-(4-cyanophenyl)-N-cyclopentyl-N-methyl-2-(5-dimethylamino-naphthalene-1-sulfonylamino)-propionamide (0.55g, 1.09 mmole). This intermediate compound was then treated according to the same procedure as Example 1 to obtain the purified title compound (0.35g, Yield : 60%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.6-7.3(m, 10H), 4.6-3.9(m, 2H), 3.11 (s, 3H), 3.0(s, 3H), 3.05-2.8(m, 2H), 2.5, 2.4(2s, 3H), 1.7-0.9(m, 10H)
 Mass(FAB, m/e) : 537($\text{M}^+ + 1$)

Example 19Synthesis of (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(5-methoxy-naphthalene-1-sulfonylamino)-propionamide

The same procedure as Preparation 3, except that 5-methoxy-1-naphthalenesulfonyl chloride was used instead of 2-naphthalenesulfonyl chloride, was carried out to obtain the intermediate (S)-3-(4-cyanophenyl)-N-cyclopentyl-N-methyl-2-(5-methoxy-naphthalene-1-sulfonylamino)-propionamide (0.18g, 0.37 mmole). This intermediate compound was then treated according to the same procedure as Example 1 to obtain the purified title compound (0.12g, Yield : 65%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.5-7.0(m, 10H), 4.6-4.0(m, 2H), 4.2(s, 3H), 3.0-2.8(m, 2H), 2.48-2.45(2s, 3H), 1.7-1.0(m, 8H)
 Mass(FAB, m/e) : 523($\text{M}^+ + 1$)

Example 20Synthesis of (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(6,7-dimethoxy-naphthalene-2-sulfonylamino)-propionamide

The same procedure as Preparation 3, except that 6,7-dimethoxy-2-naphthalenesulfonyl chloride was used instead of 2-naphthalenesulfonyl chloride, was carried out to obtain the intermediate (S)-3-(4-cyanophenyl)-N-cyclopentyl-N-methyl-2-(6,7-dimethoxy-naphthalene-2-sulfonylamino)-propionamide (2.2g, 2.2 mmole). This intermediate compound was then treated according to the same procedure as Example 1 to obtain the purified title compound (1.55g, Yield : 67%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.4-7.3(m, 9H), 4.7-4.0(m, 2H), 4.2-4.0(2s, 6H), 3.2-2.8(m, 2H), 2.6-2.2 (2s, 3H), 1.7-1.0(m, 8H)
 Mass(FAB, m/e) : 554($\text{M}^+ + 1$)

Example 21Synthesis of (S)-3-[4-(methyamidino)-phenyl]-N-cyclopentyl-N-methyl-2-(5-dimethylamino-naphthalene-1-sulfonylamino)-propionamide

The same procedure as Preparation 3, except that 5-(N,N-dimethylamino)-1-naphthalenesulfonyl chloride was used instead of 2-naphthalenesulfonyl chloride, was carried out to obtain the intermediate (S)-3-(4-cyanophenyl)-N-cyclopentyl-N-methyl-2-(5-dimethylamino-naphthalene-1-sulfonylamino)-propionamide (0.3g, 0.6 mmole). This intermediate compound was then treated according to the same procedure as Example 1, except that methylamine is used instead of 80% hydrazine hydrate, to obtain the purified title compound (0.08g, Yield : 25%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.52-7.25(m, 10H), 4.55-4.42(m, 2H), 4.23-3.98(m, 2H), 3.09(s, 3H), 3.06-2.80(m, 2H), 2.90(m, 6H), 2.45(s, 3H), 2.38(s, 3H), 1.76-0.90(m, 8H)
 Mass(FAB, m/e) : 536($\text{M}^+ + 1$)

Example 22Synthesis of (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(naphthalene-1-sulfonylamino)-propionamide

The same procedure as Preparation 3, except that 1-naphthalenesulfonyl chloride was used instead of 2-naphthalenesulfonyl chloride, was carried out to obtain the intermediate (S)-3-(4-cyanophenyl)-N-cyclopentyl-N-methyl-2-(naphthalene-1-sulfonylamino)-propionamide(0.5g, 1 mmole). This intermediate compound was then treated according to the same procedure as Example 1 to obtain the purified title compound(0.2g, Yield : 40%).

¹H NMR(CD₃OD, ppm) δ : 8.61-7.28(m, 11H), 4.55-4.41(m, 2H), 4.23-4.00(m, 2H), 3.00-2.84(m, 2H), 2.49, 2.41(2s, 3H), 1.70-1.00(m, 8H)
Mass(FAB, m/e) : 494(M⁺+1)

Example 23Synthesis of (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-[2-(naphthalen-1-yl-oxy)acetyl-amino]-propionamide

The same procedure as Preparation 6, except that (1-naphthoxy)acetic acid was used instead of 1-naphthalene acetic acid, was carried out to obtain the intermediate (S)-3-(4-cyanophenyl)-N-cyclopentyl-N-methyl-2-[2-(naphthalen-1-yloxy)-acetyl-amino]-propionamide(0.59g, 1.3 mmole). This intermediate compound was then treated according to the same procedure as Example 17 to obtain the purified title compound(0.38g, Yield : 60%).

¹H NMR(CD₃OD, ppm) δ : 8.25-6.8(m, 11H), 5.38-5.22(m, 2H), 4.77-4.39(m, 2H), 4.70(s, 2H), 3.21-3.05(m, 2H), 2.88-2.77(2s, 3H), 1.92-1.40(m, 8H)
Mass(FAB, m/e) : 488(M⁺+1)

Example 24Synthesis of (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-[2-(naphthalen-2-yl-oxy)acetyl-amino]-propionamide

The same procedure as Preparation 6, except that (2-naphthoxy)acetic acid was used instead of 1-naphthalene acetic acid, was carried out to obtain the intermediate (S)-3-[4-(amino-hydrazonomethyl)-phenyl]-N-cyclopentyl-N-methyl-2-[2-(naphthalen-2-yl-oxy)acetyl-amino]-propionamide (0.7g, 1.54 mmole). This intermediate compound was then treated according to the same procedure as Example 17 to obtain the purified title compound(0.59g, Yield : 63%).

¹H NMR(CD₃OD, ppm) δ : 7.80-7.14(m, 11H), 5.35-5.18(m, 1H), 4.76-4.35(m, 1H), 4.61(s, 2H), 3.18-3.05(m, 2H), 2.85-2.75(2s, 3H), 1.85-1.25(m, 8H)
Mass(FAB, m/e) : 488(M⁺+1)

Preparation 7Synthesis of N-t-butoxycarbonyl-N-methylaminoacetic acid methyl ester

Glycine methyl ester hydrochloride(1.0g, 8.2 mmole) was dissolved in water(12ml) and 1N aqueous sodium hydroxide solution(8.2ml), and then 1,4-dioxane(20ml) was added thereto. To this mixture was added di-t-butylidicarbonate(2.2g, 9.8 mmole) at 0°C, and the mixture was warmed to room temperature and stirred for 2 hours. The volatile substance was removed from the reaction mixture under reduced pressure and the residue was diluted with ethyl acetate, washed successively with aqueous saturated sodium hydrogen carbonate solution, dilute hydrochloric acid and saturated saline, dried over anhydrous sodium sulfate, filtered and then concentrated. The resulting product was dissolved in dimethylformamide(DMF, 10ml). To this solution was slowly added 60% sodium hydride(NaH, 0.25g, 6.4 mmole) at 0°C and then added dropwise iodomethane(CH₃I, 1.1ml). The mixture was slowly warmed to room temperature and stirred for 3 hours at the same temperature. The mixture was filtered through a Celite bed and concentrated under reduced pressure. The residue was diluted with ethyl acetate, washed successively with aqueous saturated sodium hydrogen carbonate solution, dilute hydrochloric acid and brine, dried over anhydrous sodium sulfate, filtered and then concentrated. The residue was purified by column chromatography using ethyl acetate/hexane (6/4, by volume) as an eluent to obtain the purified title compound(1.0g, Yield : 65%).

EP 0 739 886 A2

¹H NMR(CDCl₃, ppm) δ : 1.45(d, 9H), 2.95(s, 3H), 3.78(s, 3H), 3.92(s, 1H), 4.00(s, 1H)
Mass(FAB, m/e) : 204(M+1)

Preparation 8

Synthesis of [(S)-2-(t-butoxycarbonylamino)-3-(4-cyanophenyl)-propionyl]-methylamino)-acetic acid methyl ester

(S)-2-(t-butoxycarbonylamino)-3-(4-cyanophenyl)propionic acid(0.5g, 1.72 mmole) was dissolved in dimethylformamide (DMF). The resulting solution was cooled down to 0°C, and then 1-(3-dimethylaminopropyl)-3-methylcarbodiimide hydrochloride(EDC, 0.39g) and 1-hydroxybenzotriazole(HOBT, 0.28g) were added thereto and stirred until they are completely dissolved. Separately, the compound(0.35g, 1.72 mmole) prepared in Preparation 7 was dissolved in dichloromethane(2ml) and cooled down to -10°C. Trifluoroacetic acid(2ml) was added thereto and the mixture was stirred for 5 minutes, slowly warmed to room temperature, stirred again for 30 minutes and then distilled under reduced pressure to remove the volatile substance. The resulting compound thus prepared and N-methylmorpholine(1ml) were added to the solution as obtained above, and then the reaction solution was slowly warmed to room temperature and stirred for 3.5 hours. Upon completion of the reaction, the reaction solution was distilled under reduced pressure to remove the volatile substance. The residue was diluted with ethyl acetate, washed successively with aqueous saturated sodium hydrogen carbonate solution, dilute hydrochloric acid and brine, dried over anhydrous sodium sulfate, filtered and then concentrated. The residue was purified by column chromatography using ethyl acetate/hexane(3/7, by volume) as an eluent to obtain the purified title compound(0.58g, Yield : 90%).

¹H NMR(CDCl₃, ppm) δ : 1.40(m, 9H), 3.08(s, 3H), 2.95-3.25(m, 2H), 3.78(s, 3H), 3.89-4.35(m, 2H), 4.95(m, 1H), 5.52(d, 1H), 7.35(m, 2H), 7.60(m, 2H)
Mass(FAB, m/e) : 376(M+1)

Preparation 9

Synthesis of 1-[(3-(4-cyanophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl)-methylamino)-acetic acid methyl ester

The compound(0.57g, 1.52 mmole) prepared in Preparation 8 was dissolved in dichloromethane(2ml) and cooled down to -10°C, and then trifluoroacetic acid(TFA, 2ml) was added thereto. The reaction mixture was stirred for 5 minutes, slowly warmed to room temperature, stirred again for 30 minutes and then distilled under reduced pressure to remove the volatile substance. The residue was dried using a vacuum pump and then DMF(10ml) was added thereto. This solution was cooled down to -10°C and N,N-diisopropylethylamine(1ml) was added thereto. This reaction solution was warmed to room temperature and stirred for about 5 minutes and then 2-naphthalenesulfonyl chloride (0.41g, 1.82 mmole) was added thereto. The reaction mixture was stirred for one hour to complete the reaction and distilled under reduced pressure to remove the volatile substance. The residue was diluted with ethyl acetate, washed two times with water, dried over anhydrous magnesium sulfate and then filtered. The filtrate was concentrated and the residue was purified by column chromatography using ethyl acetate/hexane(1/1, by volume) as an eluent to obtain the purified title compound(0.55g, Yield : 78%).

¹H NMR(CDCl₃, ppm) δ : 2.88(s, 3H), 2.80-3.20(m, 2H), 3.80(d, 3H), 4.12(d, 2H), 4.58(m, 1H), 6.40(d, 1H), 7.20-8.40(m, 11H)
Mass(FAB, m/e) : 466(M+1)

Example 25

Synthesis of [(3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl)-methyl-amino)-acetic acid methyl ester

The compound(0.55g, 1.18 mmole) prepared in Preparation 9 was dissolved in pyridine(10ml) and the resulting solution was introduced into a branched flask, to which triethylamine(0.45ml) was added. The reaction flask was equipped so that hydrogen sulfide(H₂S) gas can be slowly introduced through one branch of the flask and discharged through another branch. The reaction solution was saturated with hydrogen sulfide gas, while stirring for about 10 minutes, during which the colorless solution was changed into green color and then gradually into dark brown. The flask was closed with rubber stopper and allowed to stand for 3 days at room temperature. Upon completion of the reaction, the reaction solution was distilled under reduced pressure to remove the volatile substance and dried by means of a vacuum pump. To the obtained yellow solid were added acetone(10ml) and iodomethane(CH₃I, 0.55ml) together and

the mixture was heated under reflux for 30 minutes. This reaction mixture was distilled again under reduced pressure to remove the volatile substance and dried by means of a vacuum pump. The residue was dissolved in absolute methanol(5ml) and then stirred. To this solution was added portionwise 80% hydrazine hydrate ($\text{H}_2\text{NNH}_2 \cdot \text{H}_2\text{O}$, 0.11ml, 1.77 mmole) over three times at an interval of 10 minutes. After the reaction is completed, the reaction solution was concentrated and then purified by HPLC to obtain the title compound(0.25g, Yield : 43%).

^1H NMR(CD_3OD , ppm) δ : 2.95(s, 3H), 2.70-3.20(m, 2H), 3.54(s, 3H), 3.80(d, 2H), 4.55(m, 1H), 7.20-8.30(m, 11H)
Mass(FAB, m/e) : 498(M+1)

Example 26

Synthesis of [(3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)propionyl]-methyl-amino}-acetic acid

The compound(160mg, 0.32 mmole) prepared in Example 25 was dissolved in the mixed solvent(4ml) of methanol and water(3:1). To this solution was slowly added lithium hydroxide hydrate ($\text{LiOH} \cdot \text{H}_2\text{O}$, 0.016g, 0.38 mmole) at 0°C and the mixture was stirred for 2 hours at room temperature. Upon completion of the reaction, the reaction solution was concentrated and purified by HPLC to obtain the title compound(50mg, Yield : 32%).

^1H NMR(CD_3OD , ppm) δ : 2.20-2.60(m, 2H), 2.48(s, 3H), 2.78(s, 3H), 2.32(m, 2H), 4.12(m, 1H), 6.80-7.80(m, 11H)
Mass(FAB, m/e) : 484(M+1)

Example 27

Synthesis of (S)-2-[(3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino)-propionic acid methyl ester

The same procedure as Preparation 7, except that (L)-alanine methyl ester was used instead of glycine methyl ester, was carried out to obtain (L)-(N-t-butoxycarbonyl-N-methyl)-alanine methyl ester, which was then treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate (S)-2-[(3-(4-cyanophenyl)-(S)-2-(naphthalene-2-sulfonylamino)propionyl]-methyl-amino)-propionic acid methyl ester(1.43g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound(0.64g, Yield : 48%).

^1H NMR(CD_3OD , ppm) δ : 0.69, 0.88(d, d, 3H), 2.79, 2.95(s, s, 3H), 2.80, 3.06(m, m, 2H), 3.48, 3.57 (s, s, 3H), 4.29(m, 1H), 4.55(m, 1H), 7.30-8.30(m, 11H)
Mass(FAB, m/e) : 512(M+1)

Example 28

Synthesis of (S)-2-[(3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino)-propionic acid

The compound prepared in Example 27 was treated according to the same procedure as Example 26 to obtain the purified title compound(0.06g, Yield : 41%).

^1H NMR(CD_3OD , ppm) δ : 0.64, 0.95(d, d, 3H), 2.76, 2.92(s, s, 3H), 2.83, 3.09(m, m, 2H), 4.37(m, 1H), 4.54(m, 1H), 7.30-8.40(m, 11H)
Mass(FAB, m/e) : 498(M+1)

Example 29

Synthesis of (R)-2-[(3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino)-propionic acid methyl ester

The same procedure as Preparation 7, except that (D)-alanine methyl ester is used instead of glycine methyl ester, was carried out to obtain (D)-(N-t-butoxycarbonyl-N-methyl)-alanine methyl ester, which was then treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate (R)-2-[(3-(4-cyanophenyl)-(S)-2-(naphthalene-

EP 0 739 886 A2

2-sulfonylamino)propionyl]-methyl-amino}-propionic acid methyl ester(0.78g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound(0.58g, Yield : 70%).

¹H NMR(CD₃OD, ppm) δ : 0.89, 1.21(d, d, 3H), 2.46, 2.94(s, s, 3H), 2.80, 3.08(m, m, 2H), 3.49, 3.78 (s, s, 3H), 4.29(m, 1H), 4.59(m, 1H), 7.30-8.40(m, 11H)
Mass(FAB, m/e) : 512(M+1)

Example 30

Synthesis of (R)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino}-propionic acid

The compound(0.03g, 0.059 mmole) prepared in Example 29 was treated according to the same procedure as Example 26 to obtain the purified title compound(0.01g, Yield : 33%).

¹H NMR(CD₃OD, ppm) δ : 0.90, 1.18(d, d, 3H), 2.44, 2.92(s, s, 3H), 2.82, 3.08(m, m, 2H), 4.33(m, 1H), 4.62(m, 1H), 7.30-8.40(m, 11H)
Mass(FAB, m/e) : 498(M+1)

Example 31

Synthesis of (R)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino]-3-methyl-butyric acid methyl ester

The same procedure as Preparation 7, except that (D)-valine methyl ester was used instead of glycine methyl ester, was carried out to obtain (D)-(N-t-butoxycarbonyl-N-methyl)-valine methyl ester, which was then treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate (R)-2-[[3-(4-cyanophenyl)-(S)-2-(naphthalene-2-sulfonylamino)propionyl]-methyl-amino]-3-methyl-butyric acid methyl ester(0.19g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound(0.11g, Yield : 55%).

¹H NMR(CD₃OD, ppm) δ : 0.59, 0.70(d, d, 3H), 0.89, 0.98(d, d, 3H), 2.09, 2.21(m, m, 1H), 2.75, 3.06 (s, s, 3H), 3.40, 3.68(s, s, 3H), 4.34, 4.38(d, d, 1H), 4.63, 4.70(m, m, 1H), 7.20-8.40(m, 11H)
Mass(FAB, m/e) : 540(M⁺+1)

Example 32

Synthesis of (R)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino]-3-methyl-butyric acid

The compound prepared in Example 31 was treated according to the same procedure as Example 26 to obtain the purified title compound(0.04g, Yield : 40%).

¹H NMR(CD₃OD, ppm) δ : 0.57, 0.63(d, d, 3H), 0.92, 0.99(d, d, 3H), 2.09, 2.18(m, m, 1H), 2.74, 3.08 (s, s, 3H), 4.18, 4.36(d, d, 1H), 4.64, 4.70(m, m, 1H), 7.20-8.40(m, 11H)
Mass(FAB, m/e) : 526(M+1)

Example 33

Synthesis of 3-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino}-propionic acid methyl ester

The same procedure as Preparation 7, except that 3-aminopropionic acid methyl ester was used instead of glycine methyl ester, was carried out to obtain 3-(N-t-butoxycarbonyl-N-methyl)-amino-propionic acid methyl ester, which was then treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate 3-[[3-(4-cyanophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino}-propionic acid methyl ester (0.69g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound(0.55g, Yield : 74%).

¹H-NMR(CD₃OD, ppm) δ : 8.31, 7.97, 7.68, 7.48(d, m, m, m, 11H), 4.62, 4.51(m, m, 1H), 3.62, 3.55 (s, s, 3H), 3.05, 2.85(m, m, 4H), 2.80, 2.45(s, s, 3H), 2.38, 1.91(m, m, 2H)
 Mass(FAB, m/e) : 512(M⁺+1)

5 Example 34

Synthesis of 3-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino]-propionic acid

10 The compound prepared in Example 33 was treated according to the same procedure as Example 26 to obtain the purified title compound(0.17g, Yield : 32%).

¹H NMR(CD₃OD, ppm) δ : 8.31, 7.98, 7.78-7.37(d, m, m, 11H), 4.65, 4.52(m, m, 1H), 3.20-2.85(m, 4H), 2.80, 2.45(s, s, 3H), 2.38, 1.91(m, m, 2H)

15 Mass(FAB, m/e) : 498(M⁺+1)

Example 35

Synthesis of 4-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino]-butyric acid methyl ester

20 The same procedure as Preparation 7, except that 4-aminobutyric acid methyl ester was used instead of glycine methyl ester, was carried out to obtain 4-(N-t-butoxycarbonyl-N-methyl)-amino-propionic acid methyl ester, which was then treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate 4-[[3-(4-cyanophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-butyric acid methyl ester (0.51g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound (0.40g, Yield : 74%).

30 ¹H NMR(CD₃OD, ppm) δ : 8.32(s, 1H), 7.98(m, 3H), 7.78-7.36(m, 7H), 4.55(m, 1H), 3.72, 3.60(s, s, 3H), 3.10, 2.81(m, m, 4H), 2.79, 2.55(s, s, 3H), 2.22(m, 1H), 1.89(m, 1H), 1.63, 1.42(m, m, 1H), 1.18(m, 1H)

Mass(FAB, m/e) : 526(M⁺+1)

Example 36

Synthesis of 4-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino]-butyric acid

The compound prepared in Example 35 was treated according to the same procedure as Example 26 to obtain the purified title compound(0.12g, Yield : 32%).

40 ¹H NMR(CD₃OD, ppm) δ : 8.32(m, 1H), 7.98(m, 3H), 7.78-7.35(m, 7H), 4.55(m, 1H), 3.05, 2.81(m, m, 4H), 2.79, 2.50(s, s, 3H), 2.18(m, 1H), 1.89 (m, 1H), 1.35(m, m, 1H), 1.16(m, 1H)

Mass(FAB, m/e) : 512(M⁺+1)

45 Preparation 10

Synthesis of N-t-butoxycarbonyl-N-cyclopropyl-acetic acid methyl ester

50 Cyclopropylamine(1.34g, 23.49 mmole) was mixed with DMF(15ml) and triethylamine(3ml) and the mixture was introduced into the reaction vessel. Methylbromoacetate(2.2ml, 23.49 mmole) and DMF(5ml) were introduced into the dropping funnel. The reaction vessel was cooled to 0°C and then the solution contained in the dropping funnel was added dropwise to the reaction vessel. Upon completion of the addition, the reaction mixture was warmed to room temperature and allowed to react for 3.5 hours. After the reaction is completed, water(10ml) and 3N sodium hydroxide(8ml) were added thereto. To the reaction mixture was added 1,4-dioxane(10ml) followed by the addition of butyloxycarbonyl anhydride(6.1g, 27.95 mmole). The reaction mixture was allowed to react for 3 hours at room temperature and distilled under reduced pressure to remove the volatile substance. The residue was diluted with ethyl acetate and washed successively with saturated sodium hydrogen carbonate, dilute hydrochloric acid and brine. The organic layer was separated, dried over anhydrous magnesium sulfate and filtered. The solvent was removed under reduced pressure from the

EP 0 739 886 A2

filtrate. The residue was purified by column chromatography (eluent : ethyl acetate/hexane (6/4, by volume)) to obtain the purified title compound (2.3g, Yield : 43%).

¹H NMR(CDCl₃, ppm) δ : 3.95(m, 2H), 3.72(m, 3H), 2.75, 2.52 (bs, bs, 1H), 1.45, 1.47(s, s, 9H), 0.80-0.45(m, 4H)
Mass(FAB, m/e) : 230(M⁺+1)

Example 37

Synthesis of [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopropylamino]-acetic acid methyl ester

The compound prepared in Preparation 10 was treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate [[3-(4-cyanophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopropylamino]-acetic acid methyl ester (0.30g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound (0.25g, Yield : 77%).

¹H NMR(CD₃OD, ppm) δ : 8.24(s, 1H), 7.93(m, 3H), 7.65(m, 3H), 7.42(d, 2H), 7.35(d, 2H), 5.02(m, 1H), 3.92(d, 1H), 3.64(d, 1H), 3.60(s, 3H), 3.19(dd, 1H), 2.80(m, 2H), 0.95, 0.85, 0.61(m, m, m, 4H)
Mass(FAB, m/e) : 524(M⁺+1)

Example 38

Synthesis of [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopropylamino]-acetic acid

The compound prepared in Example 37 was treated according to the same procedure as Example 26 to obtain the purified title compound (0.07g, Yield : 29%).

¹H NMR(CD₃OD, ppm) δ : 8.24(s, 1H), 7.93(m, 3H), 7.42(d, 2H), 7.35(d, 2H), 5.02(m, 1H), 3.95(d, 1H), 3.54(d, 1H), 3.20(dd, 1H), 2.80(m, 2H), 0.95, 0.85, 0.61(m, m, m, 4H)
Mass(FAB, m/e) : 510(M⁺+1)

Example 39

Synthesis of [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-butylamino]-acetic acid methyl ester

The same procedure as Preparation 10, except that butylamine was used instead of cyclopropylamine, was carried out to obtain N-1-butoxycarbonyl-N-butylamino-acetic acid methyl ester, which was then treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate [[3-(4-cyano-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-butylamino]-acetic acid methyl ester (0.31g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound (0.19g, Yield : 58%).

¹H NMR(CD₃OD, ppm) δ : 8.30-7.32(m, 11H), 4.32-4.09(m, 3H), 3.55(s, 3H), 3.57-2.50(m, 4H), 1.26-0.50(m, 7H)
Mass(FAB, m/e) : 540(M⁺+1)

Example 40

Synthesis of [[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-butylamino]-acetic acid

The compound prepared in Example 39 was treated according to the same procedure as Example 26 to obtain the purified title compound (0.12g, Yield : 67%).

¹H NMR(CD₃OD, ppm) δ : 8.30-7.10(m, 11H), 4.31-4.10(m, 3H), 3.52-2.55(m, 4H), 1.25-0.50(m, 7H)
Mass(FAB, m/e) : 526(M⁺+1)

Example 41Synthesis of {[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopentylamino}-acetic acid methyl ester

The same procedure as Preparation 10, except that cyclopentylamine was used instead of cyclopropylamine, was carried out to obtain N-t-butoxycarbonyl-N-cyclopentylamino-acetic acid methyl ester, which was then treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate {[3-(4-cyanophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopentylamino}-acetic acid methyl ester (0.23g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound (0.12g, Yield : 50%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.35-7.35(m, 11H), 4.66, 4.33(m, m, 1H), 4.15(m, 1H), 3.75, 3.53(m, m, 1H), 3.61(s, 3H), 3.40-2.80(m, 3H), 1.90-0.60(m, 8H)

Mass(FAB, m/e) : 552($\text{M}^+ + 1$)

Example 42Synthesis of {[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-cyclopentylamino}-acetic acid

The compound prepared in Example 41 was treated according to the same procedure as Example 26 to obtain the purified title compound (0.4g, Yield : 33%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.35-7.35(m, 1H), 4.65, 4.32(m, m, 1H), 4.15(m, 1H), 3.75, 3.52(m, m, 1H), 3.41, 3.28-3.10, 2.80(m, m, m, 3H), 1.90-0.60(m, 8H)

Mass(FAB, m/e) : 538($\text{M}^+ + 1$)

Preparation 11Synthesis of 1-(N-t-butoxycarbonyl-N-methylamino)-cyclopentane carboxylic acid methyl ester

Cycloleucine (3g, 23.2 mmole) was dissolved in 1N aqueous sodium hydroxide solution (23.2ml) and distilled water (7ml), and then 1,4-dioxane (30ml) was added thereto. To this mixture was added di-t-butylidicarbonate (6.1g, 27.8 mmole) at 0°C, and the mixture was warmed to room temperature and then stirred for 2 hours. The volatile substance was removed from the reaction mixture under reduced pressure and the residue was diluted with ethyl acetate, washed successively with aqueous saturated sodium hydrogen carbonate solution, dilute hydrochloric acid and brine, dried over anhydrous sodium sulfate, filtered and then concentrated. The resulting white solid product was dissolved in dimethylformamide (DMF, 30ml). To this solution was added potassium carbonate (4.8g, 34.8 mmole) and then added dropwise iodomethane (CH_3I , 14.4ml, 232 mmole). The reaction mixture was stirred for 2 hours at room temperature and distilled under reduced pressure to remove the volatile substance. The residual solution was diluted with ethyl acetate, washed successively with aqueous saturated sodium hydrogen carbonate solution, dilute hydrochloric acid and saturated saline, dried over anhydrous sodium sulfate, filtered and then concentrated. The resulting product was dissolved in dimethylformamide (DMF, 20ml). To this solution was slowly added 60% sodium hydride (NaH , 0.46g, 11.4 mmole) at 0°C and then added dropwise iodomethane (CH_3I , 1.8ml, 28.4 mmole). The mixture was slowly warmed to room temperature and stirred for 3 hours at the same temperature. Water was added to the reaction mixture to remove the residual sodium hydride. The mixture was filtered and concentrated under reduced pressure. The residue was diluted with ethyl acetate, washed successively with aqueous saturated sodium hydrogen carbonate solution, dilute hydrochloric acid and saturated saline, dried over anhydrous sodium sulfate, filtered and then concentrated. The residue was purified by column chromatography using ethyl acetate/hexane (3/7, by volume) as an eluent to obtain the purified title compound (2.0g, Yield : 34%).

$^1\text{H NMR}(\text{CDCl}_3, \text{ppm}) \delta$: 1.35(s, 9H), 1.62(m, 4H), 1.78(m, 2H), 2.18(m, 2H), 2.90(s, 3H), 3.62(s, 3H)

Mass(FAB, m/e) : 258($\text{M} + 1$)

Example 43

Synthesis of 1-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino]-cyclopentane carboxylic acid methyl ester

The compound prepared in Preparation 11 was treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate 1-[[3-(4-aminohydrazono-methyl)-phenyl-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid methyl ester (0.28g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound (0.56g, Yield : 53%).

¹H NMR(CD₃OD, ppm) δ : 0.52(m, 1H), 0.89(m, 1H), 1.29(m, 2H), 1.52(m, 1H), 1.76(m, 2H), 2.05(m, 1H), 2.75, 3.00(m, m, 2H), 2.88(s, 3H), 3.50(s, 3H), 4.48(m, 1H), 6.38(m, 1H), 7.30-8.40(m, 11H)
Mass(FAB, m/e) : 552(M+1)

Example 44

Synthesis of 1-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid

The compound prepared in Example 43 was treated according to the same procedure as Example 26 to obtain the purified title compound (0.01g, Yield : 17%).

¹H NMR(CD₃OD, ppm) δ : 0.42(m, 1H), 0.74(m, 1H), 1.25(m, 2H), 1.50(m, 1H), 1.78(m, 2H), 2.05(m, 1H), 2.75, 3.08(m, m, 2H), 2.98(s, 3H), 4.58(m, 1H), 7.40-8.40(m, 11H)
Mass(FAB, m/e) : 538(M+1)

Preparation 12

Synthesis of 2-(N-t-butoxycarbonyl-N-methyl)-amino-cyclopentane carboxylic acid ethyl ester

Ethyl 2-oxycyclopentane carboxylate (10ml, 67.49 mmole) was introduced together with ethanol (100ml) into a reaction vessel. Then, methylamine hydrochloride (4.69g, 68.13 mmole) and water (10ml) were added thereto to dissolve the reactants. To the reaction vessel was added sodium cyanoborohydride (4.3g, 68.43 mmole), and the mixture was adjusted to pH 6 and then allowed to react for 12 hours or more at 30 to 40°C. The reaction mixture was then concentrated under reduced pressure, cooled down to 0°C, adjusted to pH 2 using 6N hydrochloric acid and washed three times with diethyl ether. The aqueous layer was adjusted again to pH 10 and the same amount of dioxane was added thereto. To this mixture was added 1 equivalent weight of butyloxycarbonyl anhydride. This reaction mixture was allowed to react for 3 hours at room temperature. After the reaction is completed, the reaction solution was distilled under reduced pressure to remove the volatile substance, diluted with ethyl acetate and washed successively with saturated sodium hydrogen carbonate, dilute hydrochloric acid and saturated saline. The organic layer was dried over anhydrous magnesium sulfate, filtered and distilled under reduced pressure to remove the solvent. The residue was purified by column chromatography (eluent=ethyl acetate:hexane= 1:1 (v/v)) to obtain the purified title compound (5.82g, Yield : 32%).

¹H NMR(CDCl₃, ppm) δ : 4.55(m, 1H), 4.10(m, 2H), 2.79(s, 3H), 2.73(s, 1H), 2.00-1.40(m, 6H), 1.45(s, 9H), 1.24(t, 3H)
Mass(FAB, m/e) : 272(M⁺+1)

Example 45

Synthesis of 2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid ethyl ester

The compound prepared in Preparation 12 was treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate 2-[[3-(4-cyanophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid ethyl ester (0.48g). This intermediate compound was then treated according to the same procedure as Example 25 to obtain the purified title compound (0.36g, Yield : 71%).

EP 0 739 886 A2

¹H NMR(CD₃OD, ppm) δ : 8.39-7.25(m, 11H), 4.78-4.40(m, 2H), 4.05(m, 2H), 3.05(m, 1H), 2.90-2.65(m, 3H), 2.50-2.40(m, 2H), 2.05, 1.90-1.30, 0.85(m, m, m, 6H), 1.28-1.15(m, 3H)

Mass(FAB, m/e) : 566(M⁺+1)

5 Example 46

Synthesis of 2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid

10 The compound prepared in Example 45 was treated according to the same procedure as Example 26 to obtain the purified title compound(0.086g, Yield : 25%).

¹H NMR(CD₃OD, ppm) δ : 8.39-7.20(m, 11H), 4.78-4.50(m, m, 2H), 3.05(m, 1H), 2.90-2.40(m, 5H), 2.10, 1.90-1.20, 0.75(m, m, m, 6H)

15 Mass(FAB, m/e) : 538(M⁺+1)

Example 47

20 Synthesis of (S)-2-[[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methyl-amino]-3-methyl-butyrac acid methyl ester

The same procedure as Preparation 7, except that (L)-valine methyl ester is used instead of glycine methyl ester, was carried out to obtain (L)-(N-t-butoxycarbonyl-N-methyl)-valine methyl ester, which was then treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate (S)-2-[[3-(4-cyano-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propion-yl]-methylamino]-3-methyl-butyrac acid methyl ester(0.13g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound(0.09g, Yield : 69%).

¹H NMR(CD₃OD, ppm) δ : 8.30, 7.95, 7.75-7.20(m, m, m, 11H), 4.75-4.25(m, 2H), 4.65, 3.39(m, m, 3H), 3.15-2.65(m, 4H), 2.22, 2.15(s, s, 1H), 2.08, 1.90(m, m, 1H), 1.00-0.55, 0.20 (m, m, 6H)

30 Mass(FAB, m/e) : 540(M⁺+1)

Example 48

35 Synthesis of 1-[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-piperidine-(R)-2-carboxylic acid methyl ester

The same procedure as Preparation 7, except that (D)-pipecolic acid methyl ester was used instead of glycine methyl ester, was carried out and the resulting product was then treated according to the same procedure as Preparations 8 and 9 to obtain the intermediate 1-[3-(4-cyanophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-piperidine-(R)-carboxylic acid methyl ester(0.18g). This intermediate compound was treated according to the same procedure as Example 25 to obtain the purified title compound(0.16g, Yield : 84%).

¹H NMR(CD₃OD, ppm) δ : 8.32, 7.95, 7.78-7.35(m, m, m, 11H), 4.71, 4.52(m, m, 1H), 3.97, 3.80(d, d, 1H), 3.73, 3.43(s, s, 3H), 3.10, 2.83, 2.39(m, m, m, 3H), 2.05(m, 1H), 1.65-1.00, 0.30(m, m, 6H)

45 Mass(FAB, m/e) : 538(M⁺+1)

Example 49

50 Synthesis of 1-[3-(4-amidrazono-phenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-piperidine-(R)-2-carboxylic acid

The compound prepared in Example 48 was treated according to the same procedure as Example 26 to obtain the purified title compound(0.03g, Yield : 19%).

55 ¹H NMR(CD₃OD, ppm) δ : 8.35, 8.00, 7.75-7.30(m, m, m, 11H), 4.50, 4.20(m, m, 1H), 3.89(m, 1H), 3.10, 2.82, 2.45(m, m, m, 3H), 2.10(m, 1H), 1.65-1.00, 0.25(m, m, 6H)

Mass(FAB, m/e) : 524(M⁺+1)

Preparation 13Synthesis of (S)-4-[2-(butyloxycarbonyl-amino)-3-(4-methylsulfonyl-piperazinyl)-3-oxo-propyl]-benzonitrile

(S)-3-(4-cyanophenyl)-2-(butyloxycarbonylamino)propionic acid(0.5g, 1.7 mmole) was dissolved in dimethylformamide(DMF, 20ml) and then cooled to 0°C. Then, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride(EDC, 0.5g) and 1-hydroxybenzotriazole hydrate(HOBT,0.3g) were added to this solution and stirred until they are completely dissolved therein. To this reaction mixture were added 1-methanesulfonyl-piperazine(0.3g) and N-methylmorpholine(0.2ml), and then the temperature was slowly elevated to room temperature. The reaction solution was stirred for 3.5 hours. After the reaction is completed, the reaction solution was distilled under reduced pressure to remove the volatile substance and the remaining solution was diluted with ethyl acetate, washed successively with aqueous saturated sodium hydrogen carbonate solution, dilute hydrochloric acid and saturated saline, dried over anhydrous sodium sulfate, filtered and then concentrated. The residue was purified with column chromatography using ethyl acetate/hexane(6/4, by volume) as an eluent to obtain the purified title compound(0.7g, Yield : 93%).

¹H NMR(CDCl₃, ppm) δ : 7.7-7.3(m, 4H), 5.3(m, 1H), 4.8(m, 1H), 3.9-3.55(m, 2H), 3.55-2.8(m, 8H), 2.7 (s, 3H), 1.5(s, 9H)

Preparation 14Synthesis of (S)-naphthalene-2-sulfonic acid [1-(4-cyanobenzyl)-2-(4-methylsulfonyl-piperazinyl)-2-oxoethyl]amide

The compound(0.7g, 1.6 mmole) prepared in Preparation 13 was dissolved in dichloromethane(3ml) and then cooled to -10°C, and trifluoroacetic acid(TFA, 3ml) was added thereto. The reaction mixture was stirred for 5 minutes, slowly warmed to room temperature, stirred for 3 minutes and then distilled under reduced pressure to remove the volatile substance. The remaining solution was dried by means of a vacuum pump and then 20ml of DMF was added thereto. The mixture was cooled to 0°C and 2-naphthalenesulfonyl chloride(0.5g) and diisopropylethylamine(0.9ml) were added thereto. This mixture was stirred until the reactants are completely dissolved. The reaction mixture was slowly warmed to room temperature and stirred for 3 hours. Upon completion of the reaction, the reaction solution was distilled under reduced pressure to remove the volatile substance. The remaining solution was diluted with chloroform, washed successively with saturated sodium hydrogen carbonate, dilute hydrochloric acid and brine, dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by column chromatography using chloroform/methanol(95/5, by volume) as an eluent to obtain the purified title compound(0.8g, Yield : 95%).

¹H NMR(CDCl₃, ppm) δ : 8.3-7.2(m, 11H), 5.8(m, 1H), 4.5(m, 1H), 3.5-3.2(m, 2H), 2.45(s, 3H), 3.1-2.3(m, 8H)

Example 50Synthesis of (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)-benzyl]-2-(4-methylsulfonyl-piperazinyl)-2-oxo-ethyl]-amide

The compound(0.4g, 0.76 mmole) prepared in Preparation 14 was dissolved in pyridine(5ml) and the resulting solution was introduced into a branched flask, to which triethylamine(0.3ml) was added. The reaction flask was equipped so that hydrogen sulfide(H₂S) gas can be slowly introduced through one branch of the flask and discharged through another branch. The reaction solution was saturated with hydrogen sulfide gas, while stirring for about 10 minutes, during which the colorless solution was changed into green color and then gradually into dark brown. The flask was closed with rubber stopper and allowed to stand for 3 days at room temperature. Then, the reaction solution was distilled under reduced pressure to remove the volatile substance and dried by means of a vacuum pump. To the obtained yellow solid were added acetone(10ml) and iodomethane(0.5ml) together and the mixture was heated under reflux for 30 minutes. This reaction mixture was distilled again under reduced pressure to remove the volatile substance and dried by means of a vacuum pump. The residue was dissolved in absolute methanol(5ml) and then stirred. To this mixture was added portionwise 80% hydrazine hydrate(0.06ml) over three times at an interval of 10 minutes. After the reaction is completed, the reaction solution was concentrated and then purified by HPLC to obtain the title compound(0.3g, Yield : 65%).

¹H NMR(CD₃OD, ppm) δ : 8.4-7.4(m, 11H), 4.6(m, 1H), 3.5-3.2(m, 2H), 2.55(s, 3H), 3.1-2.2(m, 8H)
Mass(FAB, m/e) : 559(M⁺+1)

Example 51Synthesis of (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)-benzyl-2-oxo-2-(4-ethoxycarbonyl-piperazinyl)-ethyl]amide

The same procedure as Preparation 13, except that 1-piperazine ethyl carboxylate was used instead of 1-methanesulfonylpiperazine, was carried out and the resulting product was then treated according to the same procedure as Preparation 14 to obtain the intermediate (S)-naphthalene-2-sulfonic acid [1-(4-cyanobenzyl)-2-oxo-2-(4-ethoxycarbonyl-piperazinyl)-ethyl]amide (1g, 1.9 mmole). This intermediate compound was treated according to the same procedure as Example 50 to obtain the purified title compound(0.6g, Yield : 56%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.5-7.3(m, 11H), 4.55(m, 1H), 4.05(m, 2H), 3.2-2.4(m, 10H), 1.2(m, 3H)
 Mass(FAB, m/e) : 553($\text{M}^+ + 1$)

Example 52Synthesis of (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)-benzyl-2-(4-formyl-piperazinyl)-2-oxoethyl]amide

The same procedure as Preparation 13, except that 1-piperazine ethylcarboxaldehyde was used instead of 1-methanesulfonylpiperazine, was carried out and the resulting product was then treated according to the same procedure as Preparation 14 to obtain the intermediate (S)-naphthalene-2-sulfonic acid [1-(4-cyanobenzyl)-2-(4-formyl-piperazinyl)-2-oxoethyl]amide(7g, 1.6 mmole). This intermediate compound was treated according to the same procedure as Example 50 to obtain the purified title compound(0.5g, Yield : 62%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.36-7.44(m, 11H), 4.60(m, 1H), 3.46-2.80(m, 9H), 2.55(m, 1H)
 Mass(FAB, m/e) : 509($\text{M}^+ + 1$)

Example 53Synthesis of (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)-benzyl-2-(4-ethyl-piperazinyl)-2-oxoethyl]amide

The same procedure as Preparation 13, except that 1-ethylpiperazine was used instead of 1-methanesulfonylpiperazine, was carried out and the resulting product was then treated according to the same procedure as Preparation 14 to obtain the intermediate (S)-naphthalene-2-sulfonic acid [1-(4-cyanobenzyl)-2-(4-ethyl-piperazinyl)-2-oxoethyl]amide(0.3g, 0.6 mmole). This intermediate compound was treated according to the same procedure as Example 50 to obtain the purified title compound(0.2g, Yield : 56%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.5-7.3(m, 11H), 4.6(m, 1H), 3.5-2.7(m, 10H), 2.2(s, 2H), 1.4-1.2(m, 3H)
 Mass(FAB, m/e) : 509($\text{M}^+ + 1$)

Example 54Synthesis of (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)-benzyl-2-oxo-2-(4-phenylpiperazinyl)-ethyl]amide

The same procedure as Preparation 13, except that 1-phenylpiperazine was used instead of 1-methanesulfonylpiperazine, was carried out and the resulting product was then treated according to the same procedure as Preparation 14 to obtain the intermediate (S)-naphthalene-2-sulfonic acid [1-(4-cyanobenzyl)-2-oxo-2-(4-phenyl-piperazinyl)-ethyl]amide (0.5g, 0.97 mmole). This intermediate compound was treated according to the same procedure as Example 50 to obtain the purified title compound(0.3g, Yield : 57%).

$^1\text{H NMR}(\text{CD}_3\text{OD}, \text{ppm}) \delta$: 8.35-7.54(m, 11H), 7.20(m, 2H), 6.86(m, 1H), 6.67(m, 2H), 4.56(m, 1H), 3.45(m, 1H), 3.25-2.92(m, 5H), 3.72(m, 2H), 2.42(m, 1H), 2.05(m, 1H)
 Mass(FAB, m/e) : 557($\text{M}^+ + 1$)

Example 55Synthesis of (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)-benzyl-2-oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazinyl]-ethyl]-amide

The same procedure as Preparation 13, except that 1-(α,α,α -trifluoro-m-tolyl)piperazine was used instead of 1-methanesulfonylpiperazine, was carried out and the resulting product was then treated according to the same procedure as Preparation 14 to obtain the intermediate (S)-naphthalene-2-sulfonic acid [1-(4-cyanobenzyl)-2-oxo-2-[4-(3-trifluoromethyl-phenyl)-piperazinyl]ethyl]amide (0.5g, 0.8 mmole). This intermediate compound was treated according to the same procedure as Example 50 to obtain the purified title compound (0.3g, Yield : 55%).

^1H NMR(CD_3OD , ppm) δ : 8.35-6.7(m, 15H), 3.47(m, 1H), 3.3-3.0(m, 5H), 3.76(m, 2H), 2.45(m, 1H), 2.03(m, 1H)
Mass(FAB, m/e) : 625(M^++1)

Example 56Synthesis of (S)-naphthalene-2-sulfonic acid [2-(4-acetyl-piperazinyl)-1-(4-amidrazono)benzyl-2-oxoethyl]amide

The same procedure as Preparation 13, except that 1-acetylpiperazine was used instead of 1-methanesulfonylpiperazine, was carried out and the resulting product was then treated according to the same procedure as Preparation 14 to obtain the intermediate (S)-naphthalene-2-sulfonic acid [2-(4-acetyl-piperazinyl)-1-(4-cyanobenzyl)-2-oxo-ethyl]amide (1.25g, 2.55 mmole). This intermediate compound was treated according to the same procedure as Example 50 to obtain the purified title compound (0.7g, Yield : 53%).

^1H NMR(CD_3OD , ppm) δ : 8.4-7.4(m, 11H), 4.5(m, 1H), 3.5-3.2(m, 2H), 3.1-2.2(m, 8H), 2.0(s, 3H)
Mass(FAB, m/e) : 523(M^++1)

Example 57Synthesis of (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)-benzyl-2-oxo-2-[4-(2-hydroxyethyl)-piperazinyl]-ethyl]amide

The same procedure as Preparation 13, except that 1-piperazine ethane is used instead of 1-methanesulfonylpiperazine, was carried out and the resulting product was then treated according to the same procedure as Preparation 14 to obtain the intermediate (S)-naphthalene-2-sulfonic acid [1-(4-cyanobenzyl)-2-oxo-2-[4-(2-hydroxyethyl)-piperazinyl]ethyl]amide (0.07g, 0.14 mmole). This intermediate compound was treated according to the same procedure as Example 50 to obtain the purified title compound (0.05g, Yield : 68.5%).

^1H NMR(CD_3OD , ppm) δ : 8.35-7.30(m, 11H), 4.60(m, 1H), 3.96-3.7(m, 4H), 3.70-2.80(m, 10H)
Mass(FAB, m/e) : 525(M^++1)

Example 58Synthesis of (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)-benzyl-2-oxo-2-[4-(2-ethoxyphenyl)-piperazinyl]-ethyl]amide

The same procedure as Preparation 13, except that 1-(2-ethoxyphenyl)piperazine was used instead of 1-methanesulfonylpiperazine, was carried out and the resulting product was then treated according to the same procedure as Preparation 14 to obtain the intermediate (S)-naphthalene-2-sulfonic acid [1-(4-cyanobenzyl)-2-oxo-2-[4-(2-ethoxyphenyl)-piperazinyl]ethyl]amide (0.4g, 0.7 mmole). This intermediate compound was treated according to the same procedure as Example 50 to obtain the purified title compound (0.26g, Yield : 62%).

^1H NMR(CD_3OD , ppm) δ : 8.36-6.54(m, 15H), 4.6(m, 1H), 4.01(m, 2H), 3.48(m, 1H), 3.28-3.08(m, 5H), 2.69(m, 2H), 2.38(m, 1H), 2.08(m, 1H), 1.35(m, 3H)
Mass(FAB, m/e) : 601(M^++1)

Test 1 : Inhibitory activity for thrombin

The ability of the compound according to the present invention to inhibit the thrombin activity was determined as described in the following.

1160 μ l of 0.1M tris buffered solution (pH 7.8) containing 150mM NaCl and 0.1% PEG 8000 (polyethylene glycol, molecular weight about 8,000) was added to a 1.5ml cuvette. Chromozym TH was dissolved in dimethylsulfoxide(DMSO) in the concentration of 10mM and the resulting solution was diluted with the said tris buffered solution to the concentration of 0.1mM, which was then used as the substrate solution. 225 μ l of 0.1mM substrate solution thus prepared was added to the cuvette. The inhibitor solution was prepared by dissolving the thrombin inhibitor compound according to the present invention in dimethylsulfoxide in the concentration of 10mg/ml and then diluting the resulting solution with the said tris buffered solution to the concentration of 0.1mg/ml, 0.01mg/ml, 0.001mg/ml and 0.0001mg/ml. The resulting inhibitor solution was taken in an amount corresponding to 0 to 10 μ g as the inhibitor and then diluted with tris buffered solution to make the total volume of 100 μ l which was added to the cuvette.

15 μ l of bovine thrombin solution dissolved in above tris buffered solution in the concentration of 0.1mg/ml was added to the cuvette to initiate the enzyme hydrolysis reaction. The amount of para-nitroaniline produced for 2 minutes from the moment the enzyme is added was monitored by measuring the absorbance at 381nm. The continuous spectrum of absorbance to reaction time was depicted. The same experiments were conducted for various concentration of the inhibitor to obtain the continuous spectrum.

In each spectrum, the initial velocity V_i was obtained from the slope within initial 30 seconds of the reaction time and then a graph for the reciprocal value of initial velocity ($1/V_i$) to the inhibitor concentration was depicted. From the graph, the primary equation satisfying the points depicted thereon was obtained and then the value K_i was calculated from the x intercept of the primary equation using the enzyme reaction equation. The value K_m used for this calculation was 8.3 μ M which was obtained by changing the substrate concentration at the constant enzyme concentration.

The velocity constant K_s was obtained using the same solution in the same concentration as in determination of the value K_i but using the following experimental procedure.

Specifically, 1160 μ l of the buffered solution was added to a 1.5ml cuvette, and 15 μ l of bovine thrombin solution having 0.1mg/ml concentration and 100 μ l of the inhibitor solution were added thereto. The mixture was allowed to stand at room temperature for 15 minutes. Then, while adding 225 μ l of 0.1mM substrate solution, the change of absorbance in the course of time was monitored for 2 minutes. From the obtained continuous spectrum, the slope at the straight portion was determined and represented as the value V_s . The same experiment was conducted for various concentration of the inhibitor to obtain the value V_s at each concentration of the inhibitor from which the graph for $1/V_s$ against the inhibitor concentration was depicted. From the graph, the primary equation satisfying the points depicted thereon was obtained and then the value K_s was calculated from the x intercept of the primary equation using the enzyme reaction equation.

Separately, the inhibitory activity of the compound according to the present invention for trypsin was determined in the same manner as in determination of the inhibitory activity for thrombin as described above.

As the substrate 20 μ M solution of N-benzoyl-Val-Gly-Arg-p-nitroanilide hydrochloride was used, and the inhibitor was used in various concentrations within the range of 0 to 120 μ g. In addition, trypsin was dissolved in 0.1N HCl, adjusted to the concentration of 45 μ g/ml with the said tris buffered solution just before the experiment, and then used in an amount of 40 μ l. As in the experiment for thrombin, a total volume of the reaction solution was 1.5ml and the remaining procedure was conducted in the same manner. The value K_m used in calculation of the value K_i was determined according to the same method as in the experiment for thrombin above and was 20.2 μ M.

The inhibitory activity of the compound according to the present invention for the activity of each enzyme as determined according to the above method was represented by the values K_i and K_s and the selectivity for thrombin was represented by trypsin inhibitory activity/thrombin inhibitory activity. The result thus obtained is described in the following Table 1.

Table 1. Inhibitory activity of the compound according to the present invention for thrombin and trypsin

Comp. No. (Ex.)	Inhibitory activity for thrombin	Inhibitory activity for trypsin	Selectivity (trypsin/thrombin)
1	Ki = 0.0038 μ M Ks = 0.0011 μ M	Ki = 3.19 μ M	2900
2	Ki = 0.041 μ M		
3	Ki = 0.093 μ M		
4	Ki = 0.345 μ M		
5	Ki = 0.231 μ M		
6	Ki = 0.00435 μ M		
7	Ki = 0.0138 μ M	Ki = 363 μ M	26304
9	Ki = 0.227 μ M		
10	Ki = 0.367 μ M		
13	Ki = 0.216 μ M		
15	Ki = 0.152 μ M		
16	Ki = 0.247 μ M		
17	Ki = 0.016 μ M		
18	Ki = 0.011 μ M Ks = 0.002 μ M	Ki = 3.98 μ M	2000
19	Ki = 0.0053 μ M	Ki = 5.25 μ M	990
20	Ki = 0.0217 μ M	Ki = 2.3 μ M	106
21	Ki = 0.025 μ M Ks = 0.009 μ M	Ki > 100 μ M	>10000
22	Ki = 0.017 μ M		
23	Ki = 2.59 μ M	Ki = 21.7 μ M	8
24	Ki = 20.1 μ M		

Table 1. (continued)

Comp. No. (Ex.)	Inhibitory activity for thrombin	Inhibitory activity for trypsin	Selectivity (trypsin/ thrombin)
25	Ki = 0.259 μ M		
27	Ki = 0.065 μ M		
29	Ki = 0.089 μ M		
31	Ki = 0.165 μ M		
33	Ki = 0.035 μ M		
35	Ki = 0.612 μ M		
39	Ki = 0.165 μ M		
45	Ki = 0.346 μ M		
46	Ki = 0.880 μ M		
47	Ki = 0.665 μ M		
48	Ki = 0.013 μ M		
49	Ki = 0.406 μ M		
50	Ki = 0.011 μ M Ks = 0.006 μ M.	Ki = 5.79 μ M	965
51	Ki = 3.3 μ M		
52	Ki = 0.05 μ M		
53	Ki = 0.530 μ M		
54	Ki = 5.380 μ M		
55	Ki = 10.000 μ M		
56	Ki = 0.280 μ M		
57	Ki = 0.088 μ M		
58	Ki = 9.130 μ M		

EP 0 739 886 A2

Test 2 : Pharmacokinetic test

Test method :

Male rats and dogs were fasted for 24 hours and used as the experimental animals. 1% solution (10mg/ml) of the compound of Example 1 was prepared using physiological saline and then administered to the experimental animals via intravenous and oral route. Bloods were collected from the animals at given time intervals and then immediately mixed with methanol and zinc sulfate. Finally, the upper layer of the mixture was quantitatively analyzed at ultraviolet wavelength of 231nm to measure the drug concentration in blood by HPLC.

Test result :

The drug concentration and the pharmacokinetic parameters of the compound of Example 1 following intravenous and oral administration are listed in the following Tables 2 to 7. When the compound of Example 1 was intravenously injected, it was rapidly distributed and then slowly disappeared in both rats and dogs, whereas the elimination half-life of the compound in dogs was twice longer or more than that in rats. Furthermore, the elimination half-life of the compound of Example 1 in dogs was twice longer or more than that of commercially available argatroban(40 minutes) in human being(see, Osamu et al., Pharmacology and Therapy, vol. 14, suppl. 5, 1986). Meanwhile, it could also be identified that the bioavailability of the compound of Example 1 was 15% in rats and 61% in dogs, it was administered via oral route. However, it has been reported that argatroban was not absorbed in animals and human being when it was administered via oral route.

From the above results, the compound of Example 1 showed the better pharmacokinetic characteristics than argatroban in terms of oral absorption and elimination half-life.

Table 2

Blood concentration of the compound of Example 1 following intravenous injection of 10mg/kg in rats				
Time (min.)	Blood concentration (ng/ml)			Average(+error) (ng/ml)
	Rat-1	Rat-2	Rat-3	
1	31503	23000	31964	28822(2914)
3	8105	6773	7884	7587(412)
5	3976	3066	4223	3755(352)
10	1907	1828	1928	1888(31)
20	918	898	883	900(10)
30	631	600	591	607(12)
45	355	408	398	387(16)
60	270	242	289	267(14)
90	168	166	184	173(6)
120	76	97	119	97(12)
180	87	73	78	79(4)

Table 3

Blood concentration of compound of Example 1 following oral administration of 15mg/kg in rats				
Time (min.)	Blood concentration (ng/ml)			Average(+error) (ng/ml)
	Rat-1	Rat-2	Rat-3	
5	1610	822	570	1001(313)
10	1478	686	384	850(326)
20	569	166	97	255(147)
30	232	82	113	143(46)
45	145	107	112	121(12)
60	143	123	110	125(9)
90	66	186	119	124(35)
120	34		71	53(19)
180	24	238	64	109(66)

Table 4

Pharmacokinetic parameters of the compound of Example 1 in rats	
Parameter	Average+(error)
Elimination Half-life(min.)	47 (1)
Bioavailability (%)	15 (2)

Table 5

Blood drug concentration of the compound of Example 1 following intravenous injection of 10mg/kg in dogs			
Time (min.)	Blood concentration (ng/ml)		Average(+error) (ng/ml)
	Dog-1	Dog-2	
2	29121	36648	32885(3764)
5	9082	5228	7155(1927)
10	5935	3282	4608(1327)
20	3610	1801	2706(904)
30	2433	1335	1884(549)
45	1937	859	1398(539)
60	1471	690	1081(391)
90	1269	501	885(384)
120	1090	377	733(357)
240	544	129	337(207)
300	315	107	211(104)
360	317	92	204(112)

Table 6

Blood concentration of the compound of Example 1 following oral administration of 10mg/kg in dogs			
Time (min.)	Blood concentration (ng/ml)		Average(+error) (ng/ml)
	Dog-1	Dog-2	
10	1389	15	702(687)
20	2076	17	1047(1029)
30	1942	105	1023(918)
45	1891	558	1225(667)
60	2215	1312	1763(451)
90	1273	935	1104(169)
120	774	730	752(22)
180	663	616	640(24)
240	812	318	565(247)
300	51	222	136(86)
360	118	153	135(18)


Table 7

Pharmacokinetic parameters of the compound of Example 1 in dogs	
Parameter	Average+(error)
Elimination Half-life(min.)	98 (15)
Bioavailability (%)	61 (15)

Although this invention has been described in its preferred form with a certain degree of particularity, it is appreciated by those skilled in the art that the present disclosure of the preferred form has been made only by way of example and that numerous changes in the details of the construction, combination and arrangement of parts may be resorted to without departing from the spirit and scope of the invention.


Claims

1. A compound represented by the following formula (I):



and its pharmaceutically acceptable salt, hydrate, solvate and isomer, in which


- 20 R^1 represents acetyl substituted with aryl or aryloxy, or represents sulfonyl substituted with substituted or unsubstituted aryl or N-containing heterocyclic group,
 X represents a group of formula



- 30 R^2 and R^3 independently of one another represent hydrogen; cycloalkyl substituted or unsubstituted with carboxyl or alkoxycarbonyl; arylalkyloxy; hydroxy; or lower alkyl substituted or unsubstituted with carboxyl, alkoxycarbonyl or hydroxy, or
 35 R^2 and R^3 together with nitrogen atom to which they are attached can form a piperidine group substituted with carboxyl or alkoxycarbonyl,
 R^4 represents hydrogen, lower alkyl or lower alkoxy,
 R^5 represents alkanesulfonyl; alkoxycarbonyl; alkylcarbonyl; formyl; lower alkyl; aryl substituted or unsubstituted with alkoxy or haloalkyl; or hydroxy-substituted lower alkyl; and
 40 R^6 and R^7 independently of one another represent hydrogen, lower alkyl or amino.

2. The compound of formula (I) as defined in claim 1, in which

- 45 R^1 represents acetyl substituted with naphthyl or naphthyloxy, or represents sulfonyl substituted with naphthyl or phenyl which can be substituted or unsubstituted with one to four substituents selected from the group consisting of lower alkyl, lower alkoxy and dialkylamino,
 X represents a group of formula



- 55 R^2 and R^3 independently of one another represent C_{3-6} cycloalkyl substituted or unsubstituted with carboxyl or methoxycarbonyl; benzyloxy; lower alkyl substituted or unsubstituted with carboxyl, methoxycarbonyl or hydroxy; or hydroxy, or
 R^2 and R^3 together with nitrogen atom to which they are attached can form a piperidine group substituted with carboxyl or methoxycarbonyl,

R⁴ represents hydrogen,

R⁵ represents methanesulfonyl, ethoxycarbonyl, formyl, ethyl, phenyl, methylcarbonyl, hydroxyethyl, or phenyl which can be substituted or unsubstituted with trifluoromethyl or ethoxy,


and

R⁶ and R⁷ independently of one another represent hydrogen, methyl or amino.

3. The compound of formula (I) as defined in claim 2, wherein the compound is selected from the group consisting of:

- (S)-N-cyclopentyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-butyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-propyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-(2-benzyloxyethyl)-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-butyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-ethyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-methyl-3-[4-(methylamidino)phenyl]-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-methyl-3-[4-(1,1-dimethylamidino)phenyl]-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-methyl-3-(4-amidrazonophenyl)-2-[(4-methoxy-2,3,6-trimethylbenzene)sulfonylamino]propionamide,
- (S)-N-cyclopentyl-N-hydroxy-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-(2-hydroxyethyl)-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopentyl-N-methyl-3-[4-(methylamidino)phenyl]-2-[(4-methoxy-2,3,6-trimethylbenzene)sulfonylamino]propionamide,
- (S)-N,N-dimethyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N,N-dimethyl-3-[4-(methylamidino)phenyl]-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclohexyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-N-cyclopropyl-N-methyl-3-(4-amidrazonophenyl)-2-(2-naphthylsulfonylamino)propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(2-naphthalen-1-yl-acetylamino)-propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(5-dimethylamino-naphthalene-1-sulfonylamino)-propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(5-methoxy-naphthalene-1-sulfonylamino)-propionamide,
- (S)-2-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(6,7-dimethoxy-naphthalene-2-sulfonylamino)-propionamide,
- (S)-3-[4-(methylamidino)-phenyl]-N-cyclopentyl-N-methyl-2-(5-dimethylamino-naphthalene-1-sulfonylamino)-propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-(naphthalene-1-sulfonylamino)-propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-[2-(naphthalen-1-yl-oxy)acetylamino]-propionamide,
- (S)-3-[4-(amidrazono)-phenyl]-N-cyclopentyl-N-methyl-2-[2-(naphthalen-2-yl-oxy)acetylamino]-propionamide,
- [[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-acetic acid methyl ester,
- [[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-acetic acid,
- (S)-2-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid methyl ester,
- (S)-2-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid,
- (R)-2-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid methyl ester,
- (R)-2-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid,
- (R)-2-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-3-methylbutyric acid methyl ester,
- (R)-2-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-3-methylbutyric acid,
- 3-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid methyl ester,
- 3-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-propionic acid,
- 4-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-butyric acid methyl ester,
- 4-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-butyric acid,

- • {[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino) -propionyl]-cyclopropylamino}-acetic acid methyl ester,
 - {[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino) -propionyl]-cyclopropylamino}-acetic acid,
 - {[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino) -propionyl]-butylamino}-acetic acid methyl ester,
 - {[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino) -propionyl]-butylamino}-acetic acid,
 - {[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino) -propionyl]-cyclopentylamino}-acetic acid methyl ester,
 - {[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino) -propionyl]-cyclopentylamino}-acetic acid,
 - 1-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid methyl ester,
 - 1-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid,
 - 2-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid ethyl ester,
 - 2-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-cyclopentane carboxylic acid,
 - (S)-2-[[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-methylamino]-3-methylbutyric acid methyl ester,
 - 1-[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-piperidine-(R)-2-carboxylic acid methyl ester,
 - 1-[3-(4-amidrazonophenyl)-(S)-2-(naphthalene-2-sulfonylamino)-propionyl]-piperidine-(R)-2-carboxylic acid,
 - (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-(4-methylsulfonyl-piperazinyl)-2-oxoethyl]amide,
 - (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-(4-ethoxycarbonyl-piperazinyl)-ethyl]amide,
 - (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-(4-formyl-piperazinyl)-2-oxoethyl]amide,
 - (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-(4-ethyl-piperazinyl)-2-oxoethyl]amide,
 - (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-(4-phenyl-piperazinyl)-ethyl]amide,
 - (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-[4-(3-trifluoromethylphenyl)-piperazinyl]-ethyl]amide,
 - (S)-naphthalene-2-sulfonic acid [2-(4-acetyl-piperazinyl)-1-(4-amidrazono)benzyl-2-oxoethyl]amide,
 - (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-[4-(2-hydroxyethyl)-piperazinyl]-ethyl]amide, and
 - (S)-naphthalene-2-sulfonic acid [1-(4-amidrazono)benzyl-2-oxo-2-[4-(2-ethoxyphenyl)-piperazinyl]-ethyl]amide.
4. A pharmaceutical composition for thrombin inhibition, which comprises the compound as defined in any one of claims 1 to 3 as an active ingredient together with a pharmaceutically acceptable carrier.
5. The pharmaceutical composition as defined in claim 4, wherein the composition is formulated into an oral preparation.



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 739 886 A3**

(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:
27.01.1999 Bulletin 1999/04

(43) Date of publication A2:
30.10.1996 Bulletin 1996/44

(21) Application number: 96100433.0

(22) Date of filing: 12.01.1996

(51) Int. Cl.⁶: **C07D 295/185**, C07D 241/04,
C07C 257/18, C07C 257/22,
C07C 311/19, A61K 31/155,
A61K 31/495

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE

(30) Priority: 28.04.1995 KR 9510383

(71) Applicant: LG Chemical Limited
Seoul (KR)


(72) Inventors:
• Oh, Yeong S.
Youseong-Ku, Daejeon (KR)
• Kim, Sang S.
Seo-ku, Daejeon (KR)
• Hwang, Sang Y.
Youseong-ku, Daejeon (KR)
• Yun, Mi K.
Seo-ku, Daejeon (KR)

• Hwang, Seong R.
Youseong-ku, Daejeon (KR)
• Hong, Seong W.
Daeduk-ku, Daejeon (KR)
• Lee, Yong H.
Youseong-ku, Daejeon (KR)
• Jeong, Yi N.
Youseong-ku, Daejeon (KR)
• Lee, Koo
Youseong-ku, Daejeon (KR)
• Shin, You S.
Youseong-ku, Daejeon (KR)

(74) Representative:
R.A. KUHNEN & P.A. WACKER
Patentanwalts-gesellschaft mbH
Alois-Steinecker-Strasse 22
85354 Freising (DE)


(54) **Selective thrombin inhibitors**

(57) The present invention relates to a novel selective thrombin inhibitor having the following formula (I), which is also effective by oral administration:



in which

R¹ represents acetyl substituted with aryl or aryloxy, or represents sulfonyl substituted with substituted or unsubstituted aryl or N-containing heterocyclic group,
X represents a group of formula



- R² and R³ independently of one another represent hydrogen; cycloalkyl substituted or unsubstituted with carboxyl or alkoxy-carbonyl; arylalkyloxy; hydroxy; or lower alkyl substituted or unsubstituted with carboxyl, alkoxy-carbonyl or hydroxy, or
- R² and R³ together with nitrogen atom to which they are attached can form a piperidine group substituted with carboxyl or alkoxy-carbonyl,
- R⁴ represents hydrogen, lower alkyl or lower alkoxy.
- R⁵ represents alkanesulfonyl; alkoxy-carbonyl; alkyl-carbonyl; formyl; lower alkyl; aryl substituted or unsubstituted with alkoxy or haloalkyl; or hydroxy-substituted lower alkyl, and
- R⁶ and R⁷ independently of one another represent hydrogen, lower alkyl or amino,

and to a process for preparation thereof and a pharmaceutical composition for thrombin inhibition which comprises the compound of formula (I) as an active ingredient.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 96 10 0433

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	EP 0 097 630 A (KABIVITRUM AB) 4 January 1984 * claims 1,8; table 3 *	1,4	C07D295/185 C07D241/04 C07C257/18 C07C257/22 C07C311/19 A61K31/155 A61K31/495
X	DE 28 45 941 A (VEB ARZNEIMITTELWERK DRESDEN) 10 May 1979 * page 9 - page 10 * * page 28, first and seventh compound * * claim 7 *	1,4	
Y	WO 92 16549 A (PENTAPHARM AG) 1 October 1992 * claims 1,3-5 *	1,4,5	
D,Y	WO 94 18185 A (PENTAPHARM AG) 18 August 1994 * claims 1-6,14-16 *	1,4,5	
A,D	K. HILPERT ET AL.: JOURNAL OF MEDICINAL CHEMISTRY, vol. 37, no. 23, 1994, pages 3889-3901, XP000561935 * page 3889, right-hand column, formulae; page 3892, right-hand column, Ro 46-6240 * * tables 1-4 *	1	TECHNICAL FIELDS SEARCHED (Int.Cl.6) C07D C07C
A,D	D. W. BANNER ET AL.: THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 266, no. 30, 25 October 1991, pages 20085-93, XP002085588 * page 20086, figure 1 *	1	
A,D	DE 41 15 468 A (BEHRINGWERKE AG) 12 November 1992 * claims 1,17 *	1,4	
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 24 November 1998	Examiner Hass, C
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/82 (P04C01)

THIS PAGE BLANK (USPTO)